

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



= 45 6,624,132

10247 (43) International Publication Date
10 January 2002 (10.01.2002)

PCT

(10) International Publication Number
WO 02/02727 A1

(51) International Patent Classification⁷: C11D 3/386, 3/06

(21) International Application Number: PCT/US01/20214

(22) International Filing Date: 25 June 2001 (25.06.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
09/606,478 29 June 2000 (29.06.2000) US

(71) Applicant: ECOLAB INC. [US/US]; Ecolab Center, St. Paul, MN 55102 (US).

(72) Inventors: MAN, Victor, Fuk-Pong; Apt. 207, 1410 Carling Drive, St. Paul, MN 55108 (US). LENTSCH, Steven, Eugene; 21 Orme Court, St. Paul, MN 55116 (US). KILLEEN, Yvonne, Marie; 216 East Douglas Street, South St. Paul, MN 55075 (US).

(74) Agent: SORENSEN, Andrew, D.; Ecolab Inc., 840 Sibley Memorial Highway, Mendota Heights, MN 55118 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

Cannot overcome WO w/ Statement of Claim Amendment

(54) Title: STABLE LIQUID ENZYME COMPOSITIONS WITH ENHANCED ACTIVITY

(57) Abstract: The present invention relates to a liquid enzyme cleaning composition in which the enzyme is stable at alkaline pH and in the presence of water at concentrations of at least about 50 to about 60 weight percent. In one embodiment, the composition of the inventive stabilizes the enzyme with potassium borate.

STABLE LIQUID ENZYME COMPOSITIONS WITH ENHANCED ACTIVITY

Field of the Invention

5 The present invention relates to a liquid enzyme cleaning composition in which the enzyme is stable at alkaline pH and in the presence of water at concentrations of at least about 60 weight percent. The present enzyme cleaning composition typically yields superior soil (especially protein soil) removal properties. In one embodiment, the composition of the invention stabilizes the
10 enzyme with potassium borate.

Background of the Invention

 A major challenge of detergent development for industry, restaurants, and homes is the successful removal of soils that are resistant to conventional treatment
15 and the elimination of chemicals that are not compatible with the surroundings. One such soil is protein, and one such chemical is chlorine or chlorine yielding compounds, which can be incorporated into detergent compounds or added separately to cleaning programs for protein removal. Protein soil residues, often called protein films, occur in all food processing industries, in restaurants, in
20 laundries, and in home cleaning situations.

 In the past, chlorine has been employed to degrade protein by oxidative cleavage and hydrolysis of the peptide bond, which breaks apart large protein molecules into smaller peptide chains. The conformational structure of the protein disintegrates, dramatically lowering the binding energies, and effecting desorption
25 from the surface, followed by solubilization or suspension into the cleaning solution. The use of chlorinated detergent is not without problems, such as harshness and corrosion. In addition, a new issue may force change upon both the industry, consumers, and detergent manufacturers: the growing public concern over the health and environmental impacts of chlorine and organochlorines.

30 Detergent enzymes represent an alternative to chlorine and organochlorines. Enzymes have been employed in cleaning compositions since early in the 20th

century. However, it took years of research, until the mid 1960's, before enzymes like bacterial alkaline proteases were commercially available and which had all of the minimum pH stability and soil reactivity for detergent applications. Patents issued through the 1960s related to use of enzymes for consumer laundry pre-soak or
5 wash cycle detergent compositions and consumer automatic dishwashing detergents. Early enzyme cleaning products evolved from simple powders containing alkaline protease to more complex granular compositions containing multiple enzymes to liquid compositions containing enzymes. See, for example, U.S. Pat. No. 3,451,935 to Roald et al., issued June 24, 1969 and U.S. Pat. No. 3,519,570 to McCarty issued
10 July 7, 1970.

Liquid detergent compositions containing enzymes have advantages compared to dry powder forms. Enzyme powders or granulates tended to segregate in these mechanical mixtures resulting in non-uniform, and hence undependable, product in use. In dry compositions, humidity can cause enzyme degradation. Dry
15 powdered compositions are not as conveniently suited as liquids for rapid solubility or miscibility in cold and tepid waters nor functional as direct application products to soiled surfaces. For these reasons and for expanded applications, it became desirable to have liquid enzyme compositions.

Although water is a desirable solvent for liquid cleaning compositions, there
20 are problems in formulating enzymes into aqueous compositions. Enzymes generally denature or degrade in an aqueous medium resulting in the serious reduction or complete loss of enzyme activity. This instability results from at least two mechanisms. Enzymes have three-dimensional protein structure which can be physically or chemically changed by other solution ingredients, such as surfactants
25 and builders, causing loss of catalytic effect. Alternately when protease is present in the composition, the protease will cause proteolytic digestion of the other enzymes if they are not proteases; or of itself via a process called autolysis. The prior art discloses attempts to deal with these aqueous induced enzyme stability problems by minimizing water content or altogether eliminating water from the liquid enzyme
30 containing composition. See, for example, U.S. Pat. No. 3,697,451 to Mausner et al.

issued October 10, 1972 and U.S. Pat. No. 4,753,748 to Lailem et al. issued June 28, 1988.

In order to market an aqueous enzyme composition, the enzyme must be stabilized so that it will retain its functional activity for prolonged periods of (shelf-life or storage) time. If a stabilized enzyme system is not employed, an excess of enzyme is generally required to compensate for expected loss. However, enzymes are expensive and are in fact the most costly ingredients in a commercial detergent even though they are present in relatively minor amounts. Thus, it is no surprise that various methods of stabilizing enzyme-containing, aqueous, liquid detergent compositions are described in the patent literature. There remains a need, however, for additional methods and compositions for stabilizing enzymes in cleaning compositions, particularly at high concentrations of water and alkaline pH.

Summary of the Invention

The present invention relates to a liquid enzyme cleaning composition in which the enzyme is stable at alkaline pH and in the presence of water at concentrations of at least about 60 weight percent. The enzyme cleaning composition preferably employs potassium borate to stabilize one or more enzymes at these conditions of pH and water concentration. The present composition maintains stability of the enzyme at alkaline pH, which preferably falls in the range of about 8 to about 11, preferably greater than about 9, preferably about 9 to about 10, preferably about 9.3. The present composition maintains stability of the enzyme at concentrations of water up to about 85%, preferably in the range of about 60% by weight to about 85% by weight water, preferably about 60% by weight to about 70% by weight water, preferably 62% by weight to 69-72% by weight water.

In an embodiment, the liquid enzyme cleaning composition includes a surfactant, a deterative enzyme, a boric acid salt, and at least about 60% by weight water, formulated to retain about 100% of the deterative enzyme's initial activity at ambient temperature for at least about 11 months after forming the composition. In an embodiment, the liquid enzyme cleaning composition includes a surfactant, a deterative enzyme, a potassium borate, and at least about 60% by weight water. In an

embodiment, the liquid enzyme cleaning composition includes a surfactant, a
detergent enzyme, a boric acid salt, and at least about 80% by weight water.

Potassium borate is a preferred boric acid salt in each of these embodiments.
Potassium borate is preferably present in an amount effective to provide significant
5 stabilization of the enzyme compared to compositions without potassium borate at
the same concentrations of water. Potassium borate can be present at about 10 or 15
weight percent. Preferably, after forming the present liquid enzyme cleaning
composition including potassium borate, the detergent enzyme retains about 100% of
its initial activity for at least about 11 months at ambient temperature. Preferably,
10 after forming the present liquid enzyme cleaning composition including potassium
borate, the detergent enzyme retains at least about 80% of its initial activity at 100 °F
for at least about 50 days after forming the composition. Preferably, after forming
the present liquid enzyme cleaning composition including potassium borate, the
detergent enzyme retains at least about 50% of its initial activity at 120 °F for at least
15 about 25 days after forming the composition.

The present composition can stabilize one or more of a variety of enzyme.
Detergent enzymes that can be employed in the present compositions include a
protease, an amylase, a lipase, a cellulase, a peroxidase, a gluconase, or a mixture
thereof. Preferably the detergent enzyme is a protease, an amylase, a lipase, or a
20 mixture thereof. Preferred proteases include an alkaline protease, such as a
subtilisin. Preferred amylases include an endoamylase. Preferred lipases include a
lipolase.

The composition can also include additional ingredients such as a source of
calcium ions, a polyol, a builder, a dye, or a combination thereof. Preferably, the
25 present composition includes an amphoteric surfactant, a protease, an amylase,
and/or a lipase, potassium borate, calcium chloride, propylene glycol, citric acid salt,
and a dye. Preferably these ingredients are present at about 8% by weight surfactant,
about 2% by weight protease, about 10% to about 15% by weight boric acid salt,
about 0.25% by weight calcium chloride, about 8% by weight propylene glycol,
30 about 4% to about 7% by weight citric acid salt, and about 0.02% by weight dye.

Brief Description of the Figures

Figure 1 illustrates the amount of enzyme activity remaining in enzyme cleaning compositions with time at ambient temperature for each of formulas 1-8.

Figure 2 illustrates the amount of enzyme activity remaining in enzyme
5 cleaning compositions with time at 110 °F for each of formulas 3-6.

Figure 3 illustrates the amount of enzyme activity remaining in enzyme cleaning compositions with time at 120 °F for each of formulas 3-7.

Detailed Description of the Invention

Definitions

As used herein, weight percent, percent by weight, % by weight, and the like are synonyms that refer to the concentration of a substance as the weight of that substance divided by the weight of the composition and multiplied by 100.

As used herein, boric acid salt and borate salt are used interchangeably to
15 refer to a salt such as potassium borate or another salt obtained by or that can be visualized as being obtained by neutralization of boric acid. The weight percent of a boric acid salt or borate salt in a composition of the present invention can be expressed either as the weight percent of either the negatively charged boron containing ion, e.g. the borate or boric acid moieties, or as the weight percent of the
20 entire boric acid salt, e.g. both the negatively charged moiety and the positively charged moiety. Preferably, the weight percent refers to the entire boric acid salt. Weight percents of citric acid salts, or other acid salts, can also be expressed in these ways, preferably with reference to the entire acid salt.

As used herein, basic or alkaline pH refers to pH greater than 7, preferably
25 greater than 8 and up to about 14. Preferably basic or alkaline pH is in the range of about 8 to about 11. A preferred alkaline or basic pH value is in the range of about 9 to about 10.

As used herein, ambient temperature refers to the temperature of the surroundings of the liquid enzyme cleaning composition under normal conditions for
30 storage or transportation. Although the product may be stored and transported at

temperatures in the range of about -10 °F to about 100 °F, ambient temperature preferably refers to room temperature of about 72 °F or 25 °C.

A Stabilized Enzyme Cleaning Composition

5 The present invention relates to a liquid enzyme cleaning composition that employs a boric acid salt to provide improved enzyme stability at basic pH and in the presence of concentrations of water greater than about 50 to about 60 weight percent. In particular, the present cleaning composition containing a boric acid salt provides increased stability for proteases, for amylases, for other enzymes employed
10 with proteases, and for deterative enzymes employed in the absence of proteases. Preferably, the boric acid salt is potassium borate. The boric acid salt, e.g. potassium borate, can be obtained by any of a variety of routes. For example, commercially available boric acid salt, e.g. potassium borate, can be added to the composition. Alternatively, the boric acid salt, e.g. potassium borate, can be
15 obtained by neutralizing boric acid with a base, e.g. a potassium containing base such as potassium hydroxide.

Suitable boric acid salts provide alkalinity to the stabilized enzyme cleaning solution. Such salts include alkali metal boric acid salts; amine boric acid salts, preferably alkanolamine boric acid salts; and the like; or a combination thereof.
20 Preferred boric acid salts include potassium borate, monoethanolammonium borate, diethanolammonium borate, triethanolammonium borate, and the like, or a combination thereof. Potassium borate is a more preferred boric acid salt. The boric acid salt is preferably soluble in the composition of the invention at concentrations in excess of 5% by weight, preferably up to about 20% by weight, such as about
25 10% by weight, preferably about 15% by weight.

Advantageously, potassium borate is soluble at concentrations larger than other metal boric acid salts, particularly other alkali metal boric acid salts, particularly sodium borate. Potassium borate is employed and soluble in the present enzyme cleaning compositions at concentrations up to about 20 weight percent,
30 preferably about 5 to about 20 weight percent, preferably about 15% by weight,

preferably about 10 weight percent. Preferably this high solubility is obtained at alkaline pH, such as pH about 9 to about 10.

Potassium borate provides desirable increases in enzyme stability at basic pH compared to other buffer systems suitable for maintaining a pH above about 7,
5 preferably above about 8, preferably in the range of about 8 to about 11, more preferably about 9 to about 10. Maintaining an alkaline pH provides greater cleaning power both for most surfactants present in the cleaning composition and for the deterative enzyme, particularly when the enzyme is an alkaline protease.

Potassium borate can also provide desirable increases in enzyme stability,
10 compared to other buffer systems and agents for increasing enzyme stability, as water concentration is increased. Preferably, the present potassium borate compositions provide increased stability at concentrations of water in excess of about 60 weight percent, preferably above 65 weight percent. The upper limit to the concentration of water is set only by the amounts of other desirable or useful
15 components of the enzyme cleaning composition. That is, water can make up the entirety of the composition beyond the useful or desirable surfactant, enzyme, boric acid salt, and any additional ingredients. Typically, an upper limit for the water concentration will be about 85 weight percent. Thus the concentration of water in the present stabilized enzyme cleaning composition can be, for example, from about
20 60 weight percent to about 85 weight percent water, preferably from about 60 weight percent to about 75 weight percent water, preferably 62% to 69-72% by weight water. For example, the concentration of water in the present stabilized enzyme cleaning composition can be in a range from at least about 60%, 61%, 62%, 63%, 64%, 65%, 66%, 67%, 68%, 69%, 70%, 71%, or 72% by weight water up to about
25 61%, 62%, 63%, 64%, 65%, 66%, 67%, 68%, 69%, 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, or 85% by weight water (always selecting an upper limit that is greater than or equal to the lower limit). Advantageously, water can replace other, more expensive, solvents, cosolvents, or enzyme stabilizers employed in conventional presoak or cleaning compositions.

30 In an embodiment, the present stabilized enzyme cleaning composition includes a surfactant, a deterative enzyme, a boric acid salt, and at least about 60% by

weight water. Such a formulation can, preferably, be effective to stabilize the
deterasive enzyme at about 100% of the deterasive enzyme's initial activity at ambient
temperature for at least about 11 months after forming the composition. In an
embodiment, the present stabilized enzyme cleaning composition includes a
5 surfactant, a deterasive enzyme, a potassium borate, and at least about 60% by weight
water. In another embodiment, the present stabilized enzyme cleaning composition
includes a surfactant, a deterasive enzyme, a boric acid salt, and at least about 80% by
weight water.

In each embodiment, the stabilized enzyme cleaning solution can also
10 contain other ingredients, such as a source of calcium ions, a polyol, a builder, a dye,
or a combination thereof. In a preferred embodiment, the surfactant includes an
amphoteric surfactant, the deterasive enzyme includes a protease, the boric acid salt
includes potassium borate, the source of calcium ions includes calcium chloride, the
polyol includes propylene glycol, the builder includes citric acid salt, the dye
15 includes a dye sold under the trade name Acid Green 25, or a combination of these.
In a more preferred embodiment, the composition of the invention includes about
8% by weight surfactant, about 2% by weight protease, about 10% to about 15% by
weight boric acid salt, about 0.25% by weight calcium chloride, about 8% by weight
propylene glycol, about 4 to about 7% by weight citric acid salt, and about 0.02% by
20 weight Acid Green 25.

The boric acid salt, e.g. potassium borate, in the composition of the present
invention can provide advantageous stability to the enzyme or enzymes employed,
compared to a composition lacking the boric acid salt. The composition of the
present invention can maintain stability of an enzyme and/or prevent one enzyme
25 from degrading another enzyme. For example, the present composition can reduce
protease activity in the composition before use to a level that the protease does not
unacceptably degrade another enzyme in the composition, such as an amylase. The
protease typically degrades less than about 20% of another enzyme's activity in
about 4 weeks at ambient temperature, preferably less than about 10%, less than
30 about 5%, less than about 2%, or less than about 1%.

The composition of the present invention can also enhance the activity of an enzyme. That is, the enzyme exhibits greater activity after formulation in a composition of the invention than does control enzyme formulated in a control composition or direct from the supplier.

5 The boric acid salt, e.g. potassium borate, can provide significantly greater enzyme stability at ambient temperature and at one or more temperatures above ambient, or under other conditions indicative of storage and use stability. For example, preferably, in the present composition, the deterative enzyme retains at least about 80-100% of its initial activity at ambient temperature for at least about 30 days
10 after forming the composition; the deterative enzyme retains at least about 80-100% of its initial activity at ambient temperature for at least about 50 days after forming the composition; the deterative enzyme retains at least about 80-100% of its initial activity at ambient temperature for at least about 80 days after forming the composition; and/or the deterative enzyme retains at least about 80-100% of its initial
15 activity at ambient temperature for at least about 11 months after forming the composition. Preferably, in the present composition, the deterative enzyme retains at least about 80-100% of its initial activity at 100 °F for at least about 50 days after forming the composition and/or retains at least about 50% of its initial activity at 120 °F for at least about 25 days after forming the composition.

20 Enzyme stability and activity are typically measured by methods known to those of skill in the art. For example, the activity of the enzyme can be measured with a known enzyme assay at the time the composition is formulated and then again after the composition has been exposed to desired conditions of temperature, humidity, or the like for a predetermined time. Comparing the activity obtained
25 after exposure to the activity at an earlier time or at formulation provides a measure of enzyme stability. Suitable assays for a deterative protease include assays known to those of skill in the art and employing an azocasein substrate. Suitable assays for a deterative amylase include the Phadebas® assay for determining I-amylase activity, which is known to those of skill in the art. Enzyme assays typically include some
30 error in the determination of enzyme activity, and that error can typically be as much as about 20%, or sometimes more. Thus, an enzyme that retains full activity (or

100% of its initial activity) may show as little as about 80% of that activity in an enzyme assay. Known protocols including replicate assays and statistical analysis can be employed for determining whether the activity present is equal to (within experimental error) the initial activity, or a particular fraction of that initial activity.

5 The stabilized enzyme cleaning composition of the present invention can be employed with a variety of different surfactants, enzymes, and additional ingredients to form a variety of cleaning, destaining, and sanitizing products useful for cleaning a wide variety of articles that can be cleaned or presoaked. Preferably, the composition of the invention is formulated for cleaning or presoaking utensils, dish
10 or cooking ware, laundry, textiles, food processing surfaces, and the like. The composition of the invention can be employed for cleaning, destaining, and sanitizing products for presoaks, machine ware washing, laundry and textile cleaning and destaining, carpet cleaning and destaining, cleaning-in-place (CIP) cleaning and destaining, drain cleaning, presoaks for medical and/or dental instrument cleaning,
15 and washing or presoaks for meat cutting the equipment and other food processing surfaces.

Enzymes

 The present stabilized enzyme cleaning composition of the present invention
20 preferably includes one or more enzymes, which can provide desirable activity for removal of protein-based, carbohydrate-based, or triglyceride-based stains from substrates; for cleaning, destaining, and sanitizing presoaks, such as presoaks for flatware, cups and bowls, and pots and pans; presoaks for medical and dental instruments; or presoaks for meat cutting equipment; for machine warewashing; for
25 laundry and textile cleaning and destaining; for carpet cleaning and destaining; for cleaning-in-place and destaining-in-place; for cleaning and destaining food processing surfaces and equipment; for drain cleaning; presoaks for cleaning; and the like. Although not limiting to the present invention, enzymes suitable for the stabilized enzyme cleaning compositions can act by degrading or altering one or
30 more types of soil residues encountered on a surface or textile thus removing the soil or making the soil more removable by a surfactant or other component of the

cleaning composition. Both degradation and alteration of soil residues can improve detergency by reducing the physicochemical forces which bind the soil to the surface or textile being cleaned, i.e. the soil becomes more water soluble. For example, one or more proteases can cleave complex, macromolecular protein structures present in soil residues into simpler short chain molecules which are, of themselves, more readily desorbed from surfaces, solubilized or otherwise more easily removed by
5 detergent solutions containing said proteases.

Suitable enzymes include a protease, an amylase, a lipase, a gluconase, a cellulase, a peroxidase, or a mixture thereof of any suitable origin, such as vegetable,
10 animal, bacterial, fungal or yeast origin. Preferred selections are influenced by factors such as pH-activity and/or stability optima, thermostability, and stability to active detergents, builders and the like. In this respect bacterial or fungal enzymes are preferred, such as bacterial amylases and proteases, and fungal cellulases. Preferably the enzyme is a protease, a lipase, an amylase, or a combination thereof.

15 "Detergent enzyme", as used herein, means an enzyme having a cleaning, destaining or otherwise beneficial effect as a component of a stabilized enzyme cleaning composition for laundry, textiles, warewashing, cleaning-in-place, drains, carpets, medical or dental instruments, meat cutting tools, hard surfaces, personal care, or the like. Preferred detergent enzymes include a hydrolase such as a protease,
20 an amylase, a lipase, or a combination thereof. Preferred enzymes in stabilized enzyme cleaning compositions for warewashing or cleaning-in-place include a protease, an amylase, a cellulase, a lipase, a peroxidase, or a combination thereof.
Preferred enzymes in stabilized enzyme cleaning compositions for food processing surfaces and equipment include a protease, a lipase, an amylase, a gluconase, or a
25 combination thereof. Preferred enzymes in stabilized enzyme cleaning compositions for laundry or textiles include a protease, a cellulase, a lipase, a peroxidase, or a combination thereof. Preferred enzymes in stabilized enzyme cleaning compositions for medical or dental instruments include a protease, a lipase, or a combination thereof. Preferred enzymes in stabilized enzyme cleaning compositions for carpets
30 include a protease, an amylase, or a combination thereof. Preferred enzymes in stabilized enzyme cleaning compositions for meat cutting tools include a protease, a

lipase, or a combination thereof. Preferred enzymes in stabilized enzyme cleaning compositions for hard surfaces include a protease, a lipase, an amylase, or a combination thereof. Preferred enzymes in stabilized enzyme cleaning compositions for drains include a protease, a lipase, an amylase, or a combination thereof.

5 Enzymes are normally incorporated into a stabilized enzyme cleaning composition according to the invention in an amount sufficient to yield effective cleaning during a washing or presoaking procedure. An amount effective for cleaning refers to an amount that produces a clean, sanitary, and, preferably, corrosion free appearance to the material cleaned, particularly for flatware. An
10 amount effective for cleaning also can refer to an amount that produces a cleaning, stain removal, soil removal, whitening, deodorizing, or freshness improving effect on substrates such as utensils, pots and pans, dishware, fabrics, and the like. Typically such a cleaning effect can be achieved with amounts of enzyme from about 0.1% to about 3% by weight, preferably about 1% to about 3% by weight, of
15 the stabilized enzyme cleaning composition. Higher active levels may also be desirable in highly concentrated cleaning or presoak formulations. A presoak is preferably formulated for use upon a dilution of about 1:500, or to a formulation concentration of 2000 ppm, which puts the use concentration of the enzyme at about 10 to about 30 ppm.

20 Commercial enzymes, such as alkaline proteases, are obtainable in liquid or dried form, are sold as raw aqueous solutions or in assorted purified, processed and compounded forms, and include about 2% to about 80% by weight active enzyme generally in combination with stabilizers, buffers, cofactors, impurities and inert vehicles. The actual active enzyme content depends upon the method of
25 manufacture and is not critical, assuming the stabilized enzyme cleaning composition has the desired enzymatic activity. The particular enzyme chosen for use in the process and products of this invention depends upon the conditions of final utility, including the physical product form, use pH, use temperature, and soil types to be degraded or altered. The enzyme can be chosen to provide optimum
30 activity and stability for any given set of utility conditions.

The stabilized enzyme cleaning compositions of the present invention preferably include at least a protease. The stabilized enzyme cleaning composition of the invention has further been found, surprisingly, not only to stabilize protease for a substantially extended shelf life, but also to significantly enhance protease activity toward digesting proteins and enhancing soil removal. Further, enhanced protease activity occurs in the presence of one or more additional enzymes, such as amylase, cellulase, lipase, peroxidase, endoglucanase enzymes and mixtures thereof, preferably lipase or amylase enzymes.

A valuable reference on enzymes is "Industrial Enzymes", Scott, D., in Kirk-Othmer Encyclopedia of Chemical Technology, 3rd Edition, (editors Grayson, M. and Eckroth, D.) Vol. 9, pp. 173-224, John Wiley & Sons, New York, 1980.

Protease

A protease suitable for the stabilized enzyme cleaning composition of the present invention can be derived from a plant, an animal, or a microorganism. Preferably the protease is derived from a microorganism, such as a yeast, a mold, or a bacterium. Preferred proteases include serine proteases active at alkaline pH, preferably derived from a strain of *Bacillus* such as *Bacillus subtilis* or *Bacillus licheniformis*; these preferred proteases include native and recombinant subtilisins. The protease can be purified or a component of a microbial extract, and either wild type or variant (either chemical or recombinant). A preferred protease is neither inhibited by a metal chelating agent (sequestrant) or a thiol poison nor activated by metal ions or reducing agents, has a broad substrate specificity, is inhibited by diisopropylfluorophosphate (DFP), is an endopeptidase, has a molecular weight in the range of about 20,000 to about 40,000, and is active at a pH of about 6 to about 12 and at temperatures in a range from about 20°C to about 80°C.

Examples of proteolytic enzymes which can be employed in the stabilized enzyme cleaning composition of the invention include (with trade names) Savinase®; a protease derived from *Bacillus lentus* type, such as Maxacal®, Opticlean®, Durazym®, and Properase®; a protease derived from *Bacillus licheniformis*, such as Alcalase® and Maxatase®; and a protease derived from *Bacillus amyloliquefaciens*,

such as Primase[®]. Preferred commercially available protease enzymes include those sold under the trade names Alcalase[®], Savinase[®], Primase[®], Durazym[®], or Esperase[®] by Novo Industries A/S (Denmark); those sold under the trade names Maxatase[®], Maxacal[®], or Maxapem[®] by Gist-Brocades (Netherlands); those sold under the trade names Purafect[®], Purafect OX, and Properase by Genencor International; those sold under the trade names Opticlean[®] or Optimase[®] by Solvay Enzymes; and the like. A mixture of such proteases can also be used. For example, Purafect[®] is a preferred alkaline protease (a subtilisin) for use in detergent compositions of this invention having application in lower temperature cleaning programs, from about 30°C to about 65°C; whereas, Esperase[®] is an alkaline protease of choice for higher temperature deterative solutions, from about 50°C to about 85°C. Suitable deterative proteases are described in patent publications including: GB 1,243,784, WO 9203529 A (enzyme/inhibitor system), WO 9318140 A, and WO 9425583 (recombinant trypsin-like protease) to Novo; WO 9510591 A, WO 9507791 (a protease having decreased adsorption and increased hydrolysis), WO 95/30010, WO 95/30011, WO 95/29979, to Procter & Gamble; WO 95/10615 (*Bacillus amyloliquefaciens* subtilisin) to Genencor International; EP 130,756 A (protease A); EP 303,761 A (protease B); and EP 130,756 A. A variant protease employed in the present stabilized enzyme cleaning compositions is preferably at least 80% homologous, preferably having at least 80% sequence identity, with the amino acid sequences of the proteases in these references.

In preferred embodiments of this invention, the amount of commercial alkaline protease composite present in the composition of the invention ranges from about 0.1% by weight of deterative solution to about 3% by weight, preferably about 1% to about 3% by weight, preferably about 2% by weight, of solution of the commercial enzyme product. Typical commercially available deterative enzymes include about 5-10% of active enzyme.

Whereas establishing the percentage by weight of commercial alkaline protease required is of practical convenience for manufacturing embodiments of the present teaching, variance in commercial protease concentrates and in-situ environmental additive and negative effects upon protease activity require a more

discerning analytical technique for protease assay to quantify enzyme activity and establish correlations to soil residue removal performance and to enzyme stability within the preferred embodiment; and, if a concentrate, to use-dilution solutions.

The activity of the proteases for use in the present invention are readily expressed in
5 terms of activity units -- more specifically, Kilo-Novo Protease Units (KNPU) which are azocasein assay activity units well known to the art. A more detailed discussion of the azocasein assay procedure can be found in the publication entitled "The Use of Azoalbumin as a Substrate in the Colorimetric Determination of Peptic and Tryptic Activity", Tomarelli, R.M., Charney, J., and Harding, M.L., J. Lab. Clin.
10 Chem. 34, 428 (1949).

In preferred embodiments of the present invention, the activity of proteases present in the use-solution ranges from about 1×10^{-5} KNPU/gm solution to about 4×10^{-3} KNPU/gm solution.

Naturally, mixtures of different proteolytic enzymes may be incorporated
15 into this invention. While various specific enzymes have been described above, it is to be understood that any protease which can confer the desired proteolytic activity to the composition may be used and this embodiment of this invention is not limited in any way by specific choice of proteolytic enzyme.

20 **Amylase**

An amylase suitable for the stabilized enzyme cleaning composition of the present invention can be derived from a plant, an animal, or a microorganism. Preferably the amylase is derived from a microorganism, such as a yeast, a mold, or a bacterium. Preferred amylases include those derived from a *Bacillus*, such as *B. licheniformis*, *B. amyloliquefaciens*, *B. subtilis*, or *B. stearothermophilus*. The
25 amylase can be purified or a component of a microbial extract, and either wild type or variant (either chemical or recombinant), preferably a variant that is more stable under washing or presoak conditions than a wild type amylase.

Examples of amylase enzymes that can be employed in the stabilized enzyme
30 cleaning composition of the invention include those sold under the trade name Rapidase by Gist-Brocades® (Netherlands); those sold under the trade names

Termamyl[®], Fungamyl[®] or Duramyl[®] by Novo; Purastar STL or Purastar OXAM by Genencor; and the like. Preferred commercially available amylase enzymes include the stability enhanced variant amylase sold under the trade name Duramyl[®] by Novo. A mixture of amylases can also be used.

5 Amylases suitable for the stabilized enzyme cleaning compositions of the present invention, preferably for warewashing, include: I-amylases described in WO 95/26397, PCT/DK96/00056, and GB 1,296,839 to Novo; and stability enhanced amylases described in J. Biol. Chem., 260(11):6518-6521 (1985); WO 9510603 A, WO 9509909 A and WO 9402597 to Novo; references disclosed in WO 9402597;
10 and WO 9418314 to Genencor International. A variant I-amylase employed in the present stabilized enzyme cleaning compositions is preferably at least 80% homologous, preferably having at least 80% sequence identity, with the amino acid sequences of the proteins of these references.

Preferred amylases for use in the stabilized enzyme cleaning compositions of
15 the present invention have enhanced stability compared to certain amylases, such as Termamyl[®]. Enhanced stability refers to a significant or measurable improvement in one or more of: oxidative stability, e.g., to hydrogen peroxide/tetraacetylenediamine in buffered solution at pH 9-10; thermal stability, e.g., at common wash temperatures such as about 60 °C.; and/or alkaline
20 stability, e.g., at a pH from about 8 to about 11; each compared to a suitable control amylase, such as Termamyl[®]. Stability can be measured by methods known to those of skill in the art. Preferred enhanced stability amylases for use in the stabilized enzyme cleaning compositions of the present invention have a specific activity at least 25% higher than the specific activity of Termamyl[®] at a temperature in a range
25 of 25 °C to 55 °C and at a pH in a range of about 8 to about 10. Amylase activity for such comparisons can be measured by assays known to those of skill in the art and/or commercially available, such as the Phadebas[®] I-amylase assay.

In preferred embodiments of this invention, the amount of commercial amylase present in the composition of the invention ranges from about 0.1% by
30 weight of detergent solution to about 3% by weight, preferably about 1% to about 3% by weight, preferably about 2 % by weight, of solution of the commercial

enzyme product. Typical commercially available detergent enzymes include about 0.25-5% of active amylase.

Whereas establishing the percentage by weight of amylase required is of practical convenience for manufacturing embodiments of the present teaching,
5 variance in commercial amylase concentrates and in-situ environmental additive and negative effects upon amylase activity may require a more discerning analytical technique for amylase assay to quantify enzyme activity and establish correlations to soil residue removal performance and to enzyme stability within the preferred embodiment; and, if a concentrate, to use-dilution solutions. The activity of the
10 amylases for use in the present invention can be expressed in units known to those of skill or through amylase assays known to those of skill in the art and/or commercially available, such as the Phadebas® I-amylase assay.

Naturally, mixtures of different amylase enzymes can be incorporated into this invention. While various specific enzymes have been described above, it is to
15 be understood that any amylase which can confer the desired amylase activity to the composition can be used and this embodiment of this invention is not limited in any way by specific choice of amylase enzyme.

Cellulases

20 An cellulase suitable for the stabilized enzyme cleaning composition of the present invention can be derived from a plant, an animal, or a microorganism. Preferably the cellulase is derived from a microorganism, such as a fungus or a bacterium. Preferred cellulases include those derived from a fungus, such as *Humicola insolens*, *Humicola* strain DSM1800, or a cellulase 212-producing fungus
25 belonging to the genus *Aeromonas* and those extracted from the hepatopancreas of a marine mollusk, *Dolabella Auricula Solander*. The cellulase can be purified or a component of an extract, and either wild type or variant (either chemical or recombinant).

Examples of cellulase enzymes that can be employed in the stabilized
30 enzyme cleaning composition of the invention include those sold under the trade names Carezyme® or Celluzyme® by Novo, or Cellulase by Genencor; and the like.

A mixture of cellulases can also be used. Suitable cellulases are described in patent documents including: U.S. Pat. No. 4,435,307, GB-A-2.075.028, GB-A-2.095.275, DE-OS-2.247.832, WO 9117243, and WO 9414951 A (stabilized cellulases) to Novo.

5 In preferred embodiments of this invention, the amount of commercial cellulase present in the composition of the invention ranges from about 0.1% by weight of detergent solution to about 3% by weight, preferably about 1% to about 3% by weight, of solution of the commercial enzyme product. Typical commercially available detergent enzymes include about 5-10 percent of active
10 enzyme.

Whereas establishing the percentage by weight of cellulase required is of practical convenience for manufacturing embodiments of the present teaching, variance in commercial cellulase concentrates and in-situ environmental additive and negative effects upon cellulase activity may require a more discerning analytical
15 technique for cellulase assay to quantify enzyme activity and establish correlations to soil residue removal performance and to enzyme stability within the preferred embodiment; and, if a concentrate, to use-dilution solutions. The activity of the cellulases for use in the present invention can be expressed in units known to those of skill or through cellulase assays known to those of skill in the art and/or
20 commercially available.

Naturally, mixtures of different cellulase enzymes can be incorporated into this invention. While various specific enzymes have been described above, it is to be understood that any cellulase which can confer the desired cellulase activity to the composition can be used and this embodiment of this invention is not limited in any
25 way by specific choice of cellulase enzyme.

Lipases

A lipase suitable for the stabilized enzyme cleaning composition of the present invention can be derived from a plant, an animal, or a microorganism.
30 Preferably the lipase is derived from a microorganism, such as a fungus or a bacterium. Preferred lipases include those derived from a *Pseudomonas*, such as

Pseudomonas stutzeri ATCC 19.154, or from a *Humicola*, such as *Humicola lanuginosa* (typically produced recombinantly in *Aspergillus oryzae*). The lipase can be purified or a component of an extract, and either wild type or variant (either chemical or recombinant).

5 Examples of lipase enzymes that can be employed in the stabilized enzyme cleaning composition of the invention include those sold under the trade names Lipase P "Amano" or "Amano-P" by Amano Pharmaceutical Co. Ltd., Nagoya, Japan or under the trade name Lipolase® by Novo, and the like. Other commercially available lipases that can be employed in the present compositions include Amano-
10 CES, lipases derived from *Chromobacter viscosum*, e.g. *Chromobacter viscosum* var. *lipolyticum* NRRLB 3673 from Toyo Jozo Co., Tagata, Japan; *Chromobacter viscosum* lipases from U.S. Biochemical Corp., U.S.A. and Disoynth Co., and lipases derived from *Pseudomonas gladioli* or from *Humicola lanuginosa*.

 A preferred lipase is sold under the trade name Lipolase® by Novo. Suitable
15 lipases are described in patent documents including: WO 9414951 A (stabilized lipases) to Novo, WO 9205249, RD 94359044, GB 1,372,034, Japanese Patent Application 53,20487, laid open Feb. 24, 1978 to Amano Pharmaceutical Co. Ltd., and EP 341,947.

 In preferred embodiments of this invention, the amount of commercial lipase
20 present in the composition of the invention ranges from about 0.1% by weight of detergent solution to about 3% by weight, preferably about 1% to about 3% by weight, of solution of the commercial enzyme product. Typical commercially available detergent enzymes include about 5-10 percent of active enzyme.

 Whereas establishing the percentage by weight of lipase required is of
25 practical convenience for manufacturing embodiments of the present teaching, variance in commercial lipase concentrates and in-situ environmental additive and negative effects upon lipase activity may require a more discerning analytical technique for lipase assay to quantify enzyme activity and establish correlations to soil residue removal performance and to enzyme stability within the preferred
30 embodiment; and, if a concentrate, to use-dilution solutions. The activity of the lipases for use in the present invention can be expressed in units known to those of

skill or through lipase assays known to those of skill in the art and/or commercially available.

Naturally, mixtures of different lipase enzymes can be incorporated into this invention. While various specific enzymes have been described above, it is to be
5 understood that any lipase which can confer the desired lipase activity to the composition can be used and this embodiment of this invention is not limited in any way by specific choice of lipase enzyme.

Additional Enzymes

10 Additional enzymes suitable for use in the present stabilized enzyme cleaning compositions include a cutinase, a peroxidase, a gluconase, and the like. Suitable cutinase enzymes are described in WO 8809367 A to Genencor. Known peroxidases include horseradish peroxidase, ligninase, and haloperoxidases such as chloro- or bromo-peroxidase. Peroxidases suitable for stabilized enzyme cleaning
15 compositions are disclosed in WO 89099813 A and WO 8909813 A to Novo. Peroxidase enzymes can be used in combination with oxygen sources, e.g., percarbonate, perborate, hydrogen peroxide, and the like. Additional enzymes suitable for incorporation into the present stabilized enzyme cleaning composition are disclosed in WO 9307263 A and WO 9307260 A to Genencor International, WO
20 8908694 A to Novo, and U.S. Pat. No. 3,553,139 to McCarty et al., U.S. Pat. No. 4,101,457 to Place et al., U.S. Pat. No. 4,507,219 to Hughes and U.S. Pat. No. 4,261,868 to Hora et al.

An additional enzyme, such as a cutinase or peroxidase, suitable for the stabilized enzyme cleaning composition of the present invention can be derived from
25 a plant, an animal, or a microorganism. Preferably the enzyme is derived from a microorganism. The enzyme can be purified or a component of an extract, and either wild type or variant (either chemical or recombinant). In preferred embodiments of this invention, the amount of commercial additional enzyme, such as a cutinase or peroxidase, present in the composition of the invention ranges from
30 about 0.1% by weight of detergent solution to about 3% by weight, preferably about 1% to about 3% by weight, of solution of the commercial enzyme product. Typical

commercially available deterative enzymes include about 5-10 percent of active enzyme.

Whereas establishing the percentage by weight of additional enzyme, such as a cutinase or peroxidase, required is of practical convenience for manufacturing
5 embodiments of the present teaching, variance in commercial additional enzyme concentrates and in-situ environmental additive and negative effects upon their activity may require a more discerning analytical technique for the enzyme assay to quantify enzyme activity and establish correlations to soil residue removal performance and to enzyme stability within the preferred embodiment; and, if a
10 concentrate, to use-dilution solutions. The activity of the additional enzyme, such as a cutinase or peroxidase, for use in the present invention can be expressed in units known to those of skill or through assays known to those of skill in the art and/or commercially available.

Naturally, mixtures of different additional enzymes can be incorporated into
15 this invention. While various specific enzymes have been described above, it is to be understood that any additional enzyme which can confer the desired enzyme activity to the composition can be used and this embodiment of this invention is not limited in any way by specific choice of enzyme.

20 **Enzyme Stabilizing System**

The enzyme stabilizing system of the present invention includes a boric acid salt, such as an alkali metal borate or amine (e.g. an alkanolamine) borate, preferably an alkali metal borate, more preferably potassium borate. The enzyme stabilizing system can also include other ingredients to stabilize certain enzymes or to enhance
25 or maintain the effect of the boric acid salt.

For example, the cleaning composition of the invention can include a water-soluble source of calcium and/or magnesium ions. Calcium ions are generally more effective than magnesium ions and are preferred herein if only one type of cation is being used. Typical cleaning and/or stabilized enzyme cleaning compositions,
30 especially liquids, will include from about 1 to about 30, preferably from about 2 to about 20, more preferably from about 8 to about 12 millimoles of calcium ion per

liter of finished composition, though variation is possible depending on factors including the multiplicity, type and levels of enzymes incorporated. Preferably water-soluble calcium or magnesium salts are employed, including for example calcium chloride, calcium hydroxide, calcium formate, calcium malate, calcium maleate, calcium hydroxide and calcium acetate; more generally, calcium sulfate or magnesium salts corresponding to the listed calcium salts may be used. Further increased levels of calcium and/or magnesium may of course be useful, for example for promoting the grease-cutting action of certain types of surfactant.

Stabilizing systems of certain cleaning compositions, for example warewashing stabilized enzyme cleaning compositions, may further include from 0 to about 10%, preferably from about 0.01% to about 6% by weight, of chlorine bleach scavengers, added to prevent chlorine bleach species present in many water supplies from attacking and inactivating the enzymes, especially under alkaline conditions. While chlorine levels in water may be small, typically in the range from about 0.5 ppm to about 1.75 ppm, the available chlorine in the total volume of water that comes in contact with the enzyme, for example during warewashing, can be relatively large; accordingly, enzyme stability to chlorine in-use can be problematic. Since perborate or percarbonate, which have the ability to react with chlorine bleach, may be present in certain of the instant compositions in amounts accounted for separately from the stabilizing system, the use of additional stabilizers against chlorine, may, most generally, not be essential, though improved results may be obtainable from their use.

Suitable chlorine scavenger anions are widely known and readily available, and, if used, can be salts containing ammonium cations with sulfite, bisulfite, thiosulfite, thiosulfate, iodide, etc. Antioxidants such as carbamate, ascorbate, etc., organic amines such as ethylenediaminetetracetic acid (EDTA) or alkali metal salt thereof, monoethanolamine (MEA), and mixtures thereof can likewise be used. Likewise, special enzyme inhibition systems can be incorporated such that different enzymes have maximum compatibility. Other conventional scavengers such as bisulfate, nitrate, chloride, sources of hydrogen peroxide such as sodium perborate tetrahydrate, sodium perborate monohydrate and sodium percarbonate, as well as

phosphate, condensed phosphate, acetate, benzoate, citrate, formate, lactate, malate, tartrate, salicylate, etc., and mixtures thereof can be used if desired.

In general, since the chlorine scavenger function can be performed by ingredients separately listed under better recognized functions, there is no requirement to add a separate chlorine scavenger unless a compound performing that function to the desired extent is absent from an enzyme-containing embodiment of the invention; even then, the scavenger is added only for optimum results. Moreover, the formulator will exercise a chemist's normal skill in avoiding the use of any enzyme scavenger or stabilizer which is unacceptably incompatible, as formulated, with other reactive ingredients. In relation to the use of ammonium salts, such salts can be simply admixed with the stabilized enzyme cleaning composition but are prone to adsorb water and/or liberate ammonia during storage. Accordingly, such materials, if present, are desirably protected in a particle such as that described in U.S. Pat. No. 4,652,392, Baginski et al.

Surfactant

The surfactant or surfactant admixture of the present invention can be selected from water soluble or water dispersible nonionic, semi-polar nonionic, anionic, cationic, amphoteric, or zwitterionic surface-active agents; or any combination thereof. The particular surfactant or surfactant mixture chosen for use in the process and products of this invention can depend on the conditions of final utility, including method of manufacture, physical product form, use pH, use temperature, foam control, and soil type. Surfactants incorporated into the stabilized enzyme cleaning compositions of the present invention are preferably enzyme compatible, not substrates for the enzyme, and not inhibitors or inactivators of the enzyme. For example, when proteases and amylases are employed in the present compositions, the surfactant is preferably free of peptide and glycosidic bonds. In addition, certain cationic surfactants are known in the art to decrease enzyme effectiveness.

A preferred surfactant system of the invention can be selected from amphoteric species of surface-active agents, which offer diverse and comprehensive

commercial selection, low price; and, most important, excellent deterative effect -- meaning surface wetting, soil penetration, soil removal from the surface being cleaned, and soil suspension in the detergent solution. Despite this preference the present composition can include one or more of nonionic surfactants, anionic
5 surfactants, cationic surfactants, the sub-class of nonionic entitled semi-polar nonionics, or those surface-active agents which are characterized by persistent cationic and anionic double ion behavior, thus differing from classical amphoteric, and which are classified as zwitterionic surfactants.

Generally, the concentration of surfactant or surfactant mixture useful in
10 stabilized liquid enzyme compositions of the present invention fall in the range of from about 0.5% to about 40% by weight of the composition, preferably about 2% to about 10%, preferably about 5% to about 8%. These percentages can refer to percentages of the commercially available surfactant composition, which can contain solvents, dyes, odorants, and the like in addition to the actual surfactant. In this
15 case, the percentage of the actual surfactant chemical can be less than the percentages listed. These percentages can refer to the percentage of the actual surfactant chemical.

Preferred surfactants for the compositions of the invention include amphoteric surfactants, such as dicarboxylic coconut derivative sodium salts.

20 A typical listing of the classes and species of surfactants useful herein appears in U.S. Pat. No. 3,664,961 issued May 23, 1972, to Norris.

Nonionic Surfactant

Nonionic surfactants useful in the invention are generally characterized by
25 the presence of an organic hydrophobic group and an organic hydrophilic group and are typically produced by the condensation of an organic aliphatic, alkyl aromatic or polyoxyalkylene hydrophobic compound with a hydrophilic alkaline oxide moiety which in common practice is ethylene oxide or a polyhydration product thereof, polyethylene glycol. Practically any hydrophobic compound having a hydroxyl,
30 carboxyl, amino, or amido group with a reactive hydrogen atom can be condensed with ethylene oxide, or its polyhydration adducts, or its mixtures with alkoxylenes

such as propylene oxide to form a nonionic surface-active agent. The length of the hydrophilic polyoxyalkylene moiety which is condensed with any particular hydrophobic compound can be readily adjusted to yield a water dispersible or water soluble compound having the desired degree of balance between hydrophilic and hydrophobic properties. Useful nonionic surfactants in the present invention include:

1. Block polyoxypropylene-polyoxyethylene polymeric compounds based upon propylene glycol, ethylene glycol, glycerol, trimethylolpropane, and ethylenediamine as the initiator reactive hydrogen compound. Examples of polymeric compounds made from a sequential propoxylation and ethoxylation of initiator are commercially available under the trade names Pluronic® and Tetronic® manufactured by BASF Corp.

Pluronic® compounds are difunctional (two reactive hydrogens) compounds formed by condensing ethylene oxide with a hydrophobic base formed by the addition of propylene oxide to the two hydroxyl groups of propylene glycol. This hydrophobic portion of the molecule weighs from about 1,000 to about 4,000. Ethylene oxide is then added to sandwich this hydrophobe between hydrophilic groups, controlled by length to constitute from about 10% by weight to about 80% by weight of the final molecule.

Tetronic® compounds are tetra-functional block copolymers derived from the sequential addition of propylene oxide and ethylene oxide to ethylenediamine. The molecular weight of the propylene oxide hydrotype ranges from about 500 to about 7,000; and, the hydrophile, ethylene oxide, is added to constitute from about 10% by weight to about 80% by weight of the molecule.

2. Condensation products of one mole of alkyl phenol wherein the alkyl chain, of straight chain or branched chain configuration, or of single or dual alkyl constituent, contains from about 8 to about 18 carbon atoms with from about 3 to about 50 moles of ethylene oxide. The alkyl group can, for example, be represented by diisobutylene, di-amyl, polymerized propylene, iso-octyl, nonyl, and di-nonyl. These surfactants can be polyethylene, polypropylene, and polybutylene oxide condensates of alkyl phenols. Examples of commercial compounds of this

chemistry are available on the market under the trade names Igepal® manufactured by Rhone-Poulenc and Triton® manufactured by Union Carbide.

3. Condensation products of one mole of a saturated or unsaturated, straight or branched chain alcohol having from about 6 to about 24 carbon atoms with from about 3 to about 50 moles of ethylene oxide. The alcohol moiety can consist of mixtures of alcohols in the above delineated carbon range or it can consist of an alcohol having a specific number of carbon atoms within this range. Examples of like commercial surfactant are available under the trade names Neodol® manufactured by Shell Chemical Co. and Alfonic® manufactured by Vista Chemical Co.

4. Condensation products of one mole of saturated or unsaturated, straight or branched chain carboxylic acid having from about 8 to about 18 carbon atoms with from about 6 to about 50 moles of ethylene oxide. The acid moiety can consist of mixtures of acids in the above defined carbon atoms range or it can consist of an acid having a specific number of carbon atoms within the range. Examples of commercial compounds of this chemistry are available on the market under the trade names Nopalcol® manufactured by Henkel Corporation and Lipopeg® manufactured by Lipo Chemicals, Inc.

In addition to ethoxylated carboxylic acids, commonly called polyethylene glycol esters, other alkanolic acid esters formed by reaction with glycerides, glycerin, and polyhydric (saccharide or sorbitan/sorbitol) alcohols have application in this invention for specialized embodiments, particularly indirect food additive applications. All of these ester moieties have one or more reactive hydrogen sites on their molecule which can undergo further acylation or ethylene oxide (alkoxide) addition to control the hydrophilicity of these substances. Care must be exercised when adding these fatty ester or acylated carbohydrates to compositions of the present invention containing amylase and/or lipase enzymes because of potential incompatibility.

Examples of nonionic low foaming surfactants include:

5. Compounds from (1) which are modified, essentially reversed, by adding ethylene oxide to ethylene glycol to provide a hydrophile of designated


molecular weight; and, then adding propylene oxide to obtain hydrophobic blocks on the outside (ends) of the molecule. The hydrophobic portion of the molecule weighs from about 1,000 to about 3,100 with the central hydrophile including 10% by weight to about 80% by weight of the final molecule. These reverse Pluronics® are
 5 manufactured by BASF Corporation under the trade name Pluronic® R surfactants.

Likewise, the Tetronic® R surfactants are produced by BASF Corporation by the sequential addition of ethylene oxide and propylene oxide to ethylenediamine. The hydrophobic portion of the molecule weighs from about 2,100 to about 6,700 with the central hydrophile including 10% by weight to 80% by weight of the final
 10 molecule.

6. Compounds from groups (1), (2), (3) and (4) which are modified by "capping" or "end blocking" the terminal hydroxy group or groups (of multi-functional moieties) to reduce foaming by reaction with a small hydrophobic molecule such as propylene oxide, butylene oxide, benzyl chloride; and, short chain
 15 fatty acids, alcohols or alkyl halides containing from 1 to about 5 carbon atoms; and mixtures thereof. Also included are reactants such as thionyl chloride which convert terminal hydroxy groups to a chloride group. Such modifications to the terminal hydroxy group may lead to all-block, block-heteric, heteric-block or all-heteric nonionics.

20 Additional examples of effective low foaming nonionics include:

7. The alkylphenoxypolyethoxyalkanols of U.S. Pat No. 2,903,486 issued September 8, 1959 to Brown et al. and represented by the formula



25

in which R is an alkyl group of 8 to 9 carbon atoms, A is an alkylene chain of 3 to 4 carbon atoms, n is an integer of 7 to 16, and m is an integer of 1 to 10.

The polyalkylene glycol condensates of U.S. Pat. No. 3,048,548 issued August 7, 1962 to Martin et al. having alternating hydrophilic oxyethylene chains and hydrophobic oxypropylene chains where the weight of the terminal hydrophobic chains, the weight of the middle hydrophobic unit and the weight of the linking
 5 hydrophilic units each represent about one-third of the condensate.

The defoaming nonionic surfactants disclosed in U.S. Pat. No. 3,382,178 issued May 7 1968 to Lissant et al. having the general formula $Z[(OR)_nOH]_z$ wherein Z is alkoxylatable material, R is a radical derived from an alkaline oxide which can be ethylene and propylene and n is an integer from, for example, 10 to 2,000 or more
 10 and z is an integer determined by the number of reactive oxyalkylatable groups.

The conjugated polyoxyalkylene compounds described in U.S. Pat. No. 2,677,700, issued May 4, 1954 to Jackson et al. corresponding to the formula $Y(C_3H_6O)_n(C_2H_4O)_mH$ wherein Y is the residue of organic compound having from about 1 to 6 carbon atoms and one reactive hydrogen atom, n has an average value of
 15 at least about 6.4, as determined by hydroxyl number and m has a value such that the oxyethylene portion constitutes about 10% to about 90% by weight of the molecule.

The conjugated polyoxyalkylene compounds described in U.S. Pat. No. 2,674,619, issued April 6, 1954 to Lundsted et al. having the formula $Y[(C_3H_6O)_n(C_2H_4O)_mH]_x$ wherein Y is the residue of an organic compound having
 20 from about 2 to 6 carbon atoms and containing x reactive hydrogen atoms in which x has a value of at least about 2, n has a value such that the molecular weight of the polyoxypropylene hydrophobic base is at least about 900 and m has value such that the oxyethylene content of the molecule is from about 10% to about 90% by weight. Compounds falling within the scope of the definition for Y include, for example,
 25 propylene glycol, glycerine, pentaerythritol, trimethylolpropane, ethylenediamine and the like. The oxypropylene chains optionally, but advantageously, contain small amounts of ethylene oxide and the oxyethylene chains also optionally, but advantageously, contain small amounts of propylene oxide.

Additional conjugated polyoxyalkylene surface-active agents which are
 30 advantageously used in the compositions of this invention correspond to the formula: $P[(C_3H_6O)_n(C_2H_4O)_mH]_x$ wherein P is the residue of an organic compound

having from about 8 to 18 carbon atoms and containing x reactive hydrogen atoms in which x has a value of 1 or 2, n has a value such that the molecular weight of the polyoxyethylene portion is at least about 44 and m has a value such that the oxypropylene content of the molecule is from about 10% to about 90% by weight.

5 In either case the oxypropylene chains may contain optionally, but advantageously, small amounts of ethylene oxide and the oxyethylene chains may contain also optionally, but advantageously, small amounts of propylene oxide.

8. Polyhydroxy fatty acid amide surfactants suitable for use in the present compositions include those having the structural formula R^2CONR^1Z in
10 which: R^1 is H, C_1 - C_4 hydrocarbyl, 2-hydroxy ethyl, 2-hydroxy propyl, ethoxy, propoxy group, or a mixture thereof; R^2 is a C_5 - C_{31} hydrocarbyl, which can be straight-chain; and Z is a polyhydroxyhydrocarbyl having a linear hydrocarbyl chain with at least 3 hydroxyls directly connected to the chain, or an alkoxyated derivative (preferably ethoxylated or propoxylated) thereof. Z can be derived from a reducing
15 sugar in a reductive amination reaction; such as a glycityl moiety.

9. The alkyl ethoxylate condensation products of aliphatic alcohols with from about 0 to about 25 moles of ethylene oxide are suitable for use in the present compositions. The alkyl chain of the aliphatic alcohol can either be straight or branched, primary or secondary, and generally contains from 6 to 22 carbon atoms.

20 10. The ethoxylated C_6 - C_{18} fatty alcohols and C_6 - C_{18} mixed ethoxylated and propoxylated fatty alcohols are suitable surfactants for use in the present compositions, particularly those that are water soluble. Suitable ethoxylated fatty alcohols include the C_{10} - C_{18} ethoxylated fatty alcohols with a degree of ethoxylation of from 3 to 50.

25 11. Suitable nonionic alkylpolysaccharide surfactants, particularly for use in the present compositions include those disclosed in U.S. Pat. No. 4,565,647, Llenado, issued Jan. 21, 1986. These surfactants include a hydrophobic group containing from about 6 to about 30 carbon atoms and a polysaccharide, e.g., a polyglycoside, hydrophilic group containing from about 1.3 to about 10 saccharide
30 units. Any reducing saccharide containing 5 or 6 carbon atoms can be used, e.g., glucose, galactose and galactosyl moieties can be substituted for the glucosyl

moieties. (Optionally the hydrophobic group is attached at the 2-, 3-, 4-, etc. positions thus giving a glucose or galactose as opposed to a glucoside or galactoside.) The intersaccharide bonds can be, e.g., between the one position of the additional saccharide units and the 2-, 3-, 4-, and/or 6-positions on the preceding
5 saccharide units.

12. Fatty acid amide surfactants suitable for use the present compositions include those having the formula: $R^6\text{CON}(R^7)_2$ in which R^6 is an alkyl group containing from 7 to 21 carbon atoms and each R^7 is independently hydrogen, C_1 - C_4 alkyl, C_1 - C_4 hydroxyalkyl, or $-(C_2H_4O)_xH$, where x is in the range of from 1 to 3.

10 Preferred nonionic surfactants for the compositions of the invention include alcohol alkoxylates, EO/PO block copolymers, alkylphenol alkoxylates, and the like.


The treatise Nonionic Surfactants, edited by Schick, M.J., Vol. 1 of the Surfactant Science Series, Marcel Dekker, Inc., New York, 1983 is an excellent reference on the wide variety of nonionic compounds generally employed in the
15 practice of the present invention. A typical listing of nonionic classes, and species of these surfactants, is given in U.S. Pat. No. 3,929,678 issued to Laughlin and Heuring on Dec. 30, 1975. Further examples are given in "Surface Active Agents and Detergents" (Vol. I and II by Schwartz, Perry and Berch).

20 Semi-Polar Nonionic Surfactants

The semi-polar type of nonionic surface active agents are another class of nonionic surfactant useful in compositions of the present invention. Generally, semi-polar nonionics are high foamers and foam stabilizers, which can limit their application in CIP systems. However, within compositional embodiments of this
25 invention designed for high foam cleaning methodology, semi-polar nonionics would have immediate utility. The semi-polar nonionic surfactants include the amine oxides, phosphine oxides, sulfoxides and their alkoxylated derivatives.

13. Amine oxides are tertiary amine oxides corresponding to the general formula:

30




wherein the arrow is a conventional representation of a semi-polar bond; and, R^1 , R^2 , and R^3 may be aliphatic, aromatic, heterocyclic, alicyclic, or combinations thereof.

- 5 Generally, for amine oxides of detergent interest, R^1 is an alkyl radical of from about 8 to about 24 carbon atoms; R^2 and R^3 are alkyl or hydroxyalkyl of 1-3 carbon atoms or a mixture thereof; R^2 and R^3 can be attached to each other, e.g. through an oxygen or nitrogen atom, to form a ring structure; R^4 is an alkaline or a hydroxyalkylene group containing 2 to 3 carbon atoms; and n ranges from 0 to about 20.

- 10 Useful water soluble amine oxide surfactants are selected from the coconut or tallow alkyl di-(lower alkyl) amine oxides, specific examples of which are dodecyldimethylamine oxide, tridecyldimethylamine oxide, ehtadecyldimethylamine oxide, pentadecyldimethylamine oxide, hexadecyldimethylamine oxide, heptadecyldimethylamine oxide, octadecyldimethylamine oxide,
 15 dodecyldipropylamine oxide, tetradecyldipropylamine oxide, hexadecyldipropylamine oxide, tetradecyldibutylamine oxide, octadecyldibutylamine oxide, bis(2-hydroxyethyl)dodecylamine oxide, bis(2-hydroxyethyl)-3-dodecoxy-1-hydroxypropylamine oxide, dimethyl-(2-hydroxydodecyl)amine oxide, 3,6,9-trioctadecyldimethylamine oxide and 3-
 20 dodecoxy-2-hydroxypropyldi-(2-hydroxyethyl)amine oxide.


Useful semi-polar nonionic surfactants also include the water soluble phosphine oxides having the following structure:



25

wherein the arrow is a conventional representation of a semi-polar bond; and, R¹ is an alkyl, alkenyl or hydroxyalkyl moiety ranging from 10 to about 24 carbon atoms in chain length; and, R² and R³ are each alkyl moieties separately selected from alkyl or hydroxyalkyl groups containing 1 to 3 carbon atoms.

5 Examples of useful phosphine oxides include dimethyldecylphosphine oxide, dimethyltetradecylphosphine oxide, methylethyltetradecylphosphine oxide, dimethylhexadecylphosphine oxide, diethyl-2-hydroxyoctyldecylphosphine oxide, bis(2-hydroxyethyl)dodecylphosphine oxide, and
 bis(hydroxymethyl)tetradecylphosphine oxide. Semi-polar nonionic surfactants
 10 useful herein also include the water soluble sulfoxide compounds which have the structure:



15

wherein the arrow is a conventional representation of a semi-polar bond; and, R¹ is an alkyl or hydroxyalkyl moiety of about 8 to about 28 carbon atoms, from 0 to about 5 ether linkages and from 0 to about 2 hydroxyl substituents; and R² is an alkyl moiety consisting of alkyl and hydroxyalkyl groups having 1 to 3 carbon
 20 atoms.

Useful examples of these sulfoxides include dodecyl methyl sulfoxide; 3-hydroxy tridecyl methyl sulfoxide; 3-methoxy tridecyl methyl sulfoxide; and 3-hydroxy-4-dodecoxybutyl methyl sulfoxide.

Preferred semi-polar nonionic surfactants for the compositions of the
 25 invention include dimethyl amine oxides, such as lauryl dimethyl amine oxide, myristyl dimethyl amine oxide, cetyl dimethyl amine oxide, combinations thereof, and the like.

Anionic Surfactants

Also useful in the present invention are surface active substances which are categorized as anionics because the charge on the hydrophobe is negative; or surfactants in which the hydrophobic section of the molecule carries no charge unless the pH is elevated to neutrality or above (e.g. carboxylic acids). Carboxylate, sulfonate, sulfate and phosphate are the polar (hydrophilic) solubilizing groups found in anionic surfactants. Of the cations (counter ions) associated with these polar groups, sodium, lithium and potassium impart water solubility; ammonium and substituted ammonium ions provide both water and oil solubility; and, calcium, barium, and magnesium promote oil solubility.

As those skilled in the art understand, anionics are excellent deterative surfactants and are therefore, favored additions to heavy duty detergent compositions. Generally, however, anionics have high foam profiles which limit their use alone or at high concentration levels in cleaning systems such as CIP circuits that require strict foam control. Anionics are very useful additives to preferred compositions of the present invention. Further, anionic surface active compounds are useful to impart special chemical or physical properties other than detergency within the composition. Anionics can be employed as gelling agents or as part of a gelling or thickening system. Anionics are excellent solubilizers and can be used for hydrotropic effect and cloud point control.

The majority of large volume commercial anionic surfactants can be subdivided into five major chemical classes and additional sub-groups known to those of skill in the art and described in "Surfactant Encyclopedia", Cosmetics & Toiletries, Vol. 104 (2) 71-86 (1989). The first class includes acylamino acids (and salts), such as acylgluamates, acyl peptides, sarcosinates (e.g. N-acyl sarcosinates), taurates (e.g. N-acyl taurates and fatty acid amides of methyl tauride), and the like. The second class includes carboxylic acids (and salts), such as alkanoic acids (and alkanoates), ester carboxylic acids (e.g. alkyl succinates), ether carboxylic acids, and the like. The third class includes phosphoric acid esters and their salts. The fourth class includes sulfonic acids (and salts), such as isethionates (e.g. acyl isethionates), alkylaryl sulfonates, alkyl sulfonates, sulfosuccinates (e.g. monoesters and diesters

of sulfosuccinate), and the like. The fifth class includes sulfuric acid esters (and salts), such as alkyl ether sulfates, alkyl sulfates, and the like. Although each of these classes of anionic surfactants can be employed in the present compositions, it should be noted that certain of these anionic surfactants may be incompatible with the enzymes incorporated into the present invention. For example, the acyl-amino acids and salts may be incompatible with proteolytic enzymes because of their peptide structure.

Anionic sulfate surfactants suitable for use in the present compositions include the linear and branched primary and secondary alkyl sulfates, alkyl ethoxysulfates, fatty oleyl glycerol sulfates, alkyl phenol ethylene oxide ether sulfates, the C₃-C₁₇, acyl-N-(C₁-C₄ alkyl) and -N-(C₁-C₂ hydroxyalkyl) glucamine sulfates, and sulfates of alkylpolysaccharides such as the sulfates of alkylpolyglucoside (the nonionic nonsulfated compounds being described herein).

Examples of suitable synthetic, water soluble anionic detergent compounds include the ammonium and substituted ammonium (such as mono-, di- and triethanolamine) and alkali metal (such as sodium, lithium and potassium) salts of the alkyl mononuclear aromatic sulfonates such as the alkyl benzene sulfonates containing from about 5 to about 18 carbon atoms in the alkyl group in a straight or branched chain, e.g., the salts of alkyl benzene sulfonates or of alkyl toluene, xylene, cumene and phenol sulfonates; alkyl naphthalene sulfonate, diamyl naphthalene sulfonate, and dinonyl naphthalene sulfonate and alkoxylated derivatives.

Anionic carboxylate surfactants suitable for use in the present compositions include the alkyl ethoxy carboxylates, the alkyl polyethoxy polycarboxylate surfactants and the soaps (e.g. alkyl carboxyls). Secondary soap surfactants (e.g. alkyl carboxyl surfactants) useful in the present compositions include those which contain a carboxyl unit connected to a secondary carbon. The secondary carbon can be in a ring structure, e.g. as in p-octyl benzoic acid, or as in alkyl-substituted cyclohexyl carboxylates. The secondary soap surfactants typically contain no ether linkages, no ester linkages and no hydroxyl groups. Further, they typically lack nitrogen atoms in the head-group (amphiphilic portion). Suitable secondary soap

surfactants typically contain 11-13 total carbon atoms, although more carbons atoms (e.g., up to 16) can be present.

Other anionic detergents suitable for use in the present compositions include olefin sulfonates, such as long chain alkene sulfonates, long chain hydroxyalkane sulfonates or mixtures of alkenesulfonates and hydroxyalkane-sulfonates. Also included are the alkyl sulfates, alkyl poly(ethyleneoxy) ether sulfates and aromatic poly(ethyleneoxy) sulfates such as the sulfates or condensation products of ethylene oxide and nonyl phenol (usually having 1 to 6 oxyethylene groups per molecule. Resin acids and hydrogenated resin acids are also suitable, such as rosin, hydrogenated rosin, and resin acids and hydrogenated resin acids present in or derived from tallow oil.

The particular salts will be suitably selected depending upon the particular formulation and the needs therein.

Further examples of suitable anionic surfactants are given in "Surface Active Agents and Detergents" (Vol. I and II by Schwartz, Perry and Berch). A variety of such surfactants are also generally disclosed in U.S. Pat. No. 3,929,678, issued Dec. 30, 1975 to Laughlin, et al. at Column 23, line 58 through Column 29, line 23.


Cationic Surfactants

Surface active substances are classified as cationic if the charge on the hydrotrope portion of the molecule is positive. Surfactants in which the hydrotrope carries no charge unless the pH is lowered close to neutrality or lower, but which are then cationic (e.g. alkyl amines), are also included in this group. In theory, cationic surfactants may be synthesized from any combination of elements containing an "onium" structure R_nX+Y^- and could include compounds other than nitrogen (ammonium) such as phosphorus (phosphonium) and sulfur (sulfonium). In practice, the cationic surfactant field is dominated by nitrogen containing compounds, probably because synthetic routes to nitrogenous cationics are simple and straightforward and give high yields of product, which can make them less expensive.

Cationic surfactants preferably include, more preferably refer to, compounds containing at least one long carbon chain hydrophobic group and at least one positively charged nitrogen. The long carbon chain group may be attached directly to the nitrogen atom by simple substitution; or more preferably indirectly by a bridging functional group or groups in so-called interrupted alkylamines and amido amines. Such functional groups can make the molecule more hydrophilic and/or more water dispersible, more easily water solubilized by co-surfactant mixtures, and/or water soluble. For increased water solubility, additional primary, secondary or tertiary amino groups can be introduced or the amino nitrogen can be quaternized with low molecular weight alkyl groups. Further, the nitrogen can be a part of branched or straight chain moiety of varying degrees of unsaturation or of a saturated or unsaturated heterocyclic ring. In addition, cationic surfactants may contain complex linkages having more than one cationic nitrogen atom.

The surfactant compounds classified as amine oxides, amphoterics and zwitterions are themselves typically cationic in near neutral to acidic pH solutions and can overlap surfactant classifications. Polyoxyethylated cationic surfactants generally behave like nonionic surfactants in alkaline solution and like cationic surfactants in acidic solution.

The simplest cationic amines, amine salts and quaternary ammonium compounds can be schematically drawn thus:



in which, R represents a long alkyl chain, R', R'', and R''' may be either long alkyl chains or smaller alkyl or aryl groups or hydrogen and X represents an anion. The amine salts and quaternary ammonium compounds are preferred for practical use in this invention due to their high degree of water solubility.


The majority of large volume commercial cationic surfactants can be subdivided into four major classes and additional sub-groups known to those or skill in the art and described in "Surfactant Encyclopedia", Cosmetics & Toiletries, Vol. 104 (2) 86-96 (1989). The first class includes alkylamines and their salts. The

5 second class includes alkyl imidazolines. The third class includes ethoxylated amines. The fourth class includes quaternaries, such as alkylbenzyldimethylammonium salts, alkyl benzene salts, heterocyclic ammonium salts, tetra alkylammonium salts, and the like. Cationic surfactants are known to have a variety of properties that can be beneficial in the present compositions. These






10 desirable properties can include detergency in compositions of or below neutral pH, antimicrobial efficacy, thickening or gelling in cooperation with other agents, and the like.

Cationic surfactants useful in the compositions of the present invention include those having the formula $R^1_m R^2_x Y_L Z$ wherein each R^1 is an organic group

15 containing a straight or branched alkyl or alkenyl group optionally substituted with up to three phenyl or hydroxy groups and optionally interrupted by up to four of the following structures:



20






or an isomer or mixture of these structures, and which contains from about 8 to 22 carbon atoms. The R^1 groups can additionally contain up to 12 ethoxy groups. m is a number from 1 to 3. Preferably, no more than one R^1 group in a molecule has 16 or more carbon atoms when m is 2 or more than 12 carbon atoms when m is 3. Each


5 R^2 is an alkyl or hydroxyalkyl group containing from 1 to 4 carbon atoms or a benzyl group with no more than one R^2 in a molecule being benzyl, and x is a number from 0 to 11, preferably from 0 to 6. The remainder of any carbon atom positions on the Y group are filled by hydrogens.

Y is can be a group including, but not limited to:

10








$p = \text{about } 1 \text{ to } 12$



$p = \text{about } 1 \text{ to } 12$

15





or a mixture thereof. Preferably, L is 1 or 2, with the Y groups being separated by a moiety selected from R¹ and R² analogs (preferably alkylene or alkenylene) having from 1 to about 22 carbon atoms and two free carbon single bonds when L is 2. Z is a water soluble anion, such as a halide, sulfate, methylsulfate, hydroxide, or nitrate anion, particularly preferred being chloride, bromide, iodide, sulfate or methyl sulfate anions, in a number to give electrical neutrality of the cationic component.

10

Amphoteric Surfactants

Amphoteric, or ampholytic, surfactants contain both a basic and an acidic hydrophilic group and an organic hydrophobic group. These ionic entities may be any of anionic or cationic groups described herein for other types of surfactants. A basic nitrogen and an acidic carboxylate group are the typical functional groups employed as the basic and acidic hydrophilic groups. In a few surfactants, sulfonate, sulfate, phosphonate or phosphate provide the negative charge.


Amphoteric surfactants can be broadly described as derivatives of aliphatic secondary and tertiary amines, in which the aliphatic radical may be straight chain or branched and wherein one of the aliphatic substituents contains from about 8 to 18 carbon atoms and one contains an anionic water solubilizing group, e.g., carboxy, sulfo, sulfato, phosphato, or phosphono. Amphoteric surfactants are subdivided into two major classes known to those of skill in the art and described in "Surfactant Encyclopedia" Cosmetics & Toiletries, Vol. 104 (2) 69-71 (1989). The first class includes acyl/dialkyl ethylenediamine derivatives (e.g. 2-alkyl hydroxyethyl imidazoline derivatives) and their salts. The second class includes N-alkylamino

25

acids and their salts. Some amphoteric surfactants can be envisioned as fitting into both classes.

Amphoteric surfactants can be synthesized by methods known to those of skill in the art. For example, 2-alkyl hydroxyethyl imidazoline is synthesized by condensation and ring closure of a long chain carboxylic acid (or a derivative) with dialkyl ethylenediamine. Commercial amphoteric surfactants are derivatized by subsequent hydrolysis and ring-opening of the imidazoline ring by alkylation -- for example with chloroacetic acid or ethyl acetate. During alkylation, one or two carboxy-alkyl groups react to form a tertiary amine and an ether linkage with differing alkylating agents yielding different tertiary amines.

Long chain imidazole derivatives having application in the present invention generally have the general formula:



15

wherein R is an acyclic hydrophobic group containing from about 8 to 18 carbon atoms and M is a cation to neutralize the charge of the anion, generally sodium.

Commercially prominent imidazoline-derived amphoterics that can be employed in the present compositions include for example: Cocoamphopropionate,

20 Cocoamphocarboxy-propionate, Cocoamphoglycinate, Cocoamphocarboxy-glycinate, Cocoamphopropyl-sulfonate, and Cocoamphocarboxy-propionic acid.

Preferred amphocarboxylic acids are produced from fatty imidazolines in which the dicarboxylic acid functionality of the amphodicarboxylic acid is diacetic acid and/or dipropionic acid.

The carboxymethylated compounds (glycinates) described herein above frequently are called betaines. Betaines are a special class of amphoteric discussed herein below in the section entitled, Zwitterion Surfactants.

Long chain N-alkylamino acids are readily prepared by reaction RNH_2 , in which $\text{R}=\text{C}_8\text{-C}_{18}$ straight or branched chain alkyl, fatty amines with halogenated carboxylic acids. Alkylation of the primary amino groups of an amino acid leads to secondary and tertiary amines. Alkyl substituents may have additional amino groups that provide more than one reactive nitrogen center. Most commercial N-alkylamine acids are alkyl derivatives of beta-alanine or beta-N(2-carboxyethyl) alanine. Examples of commercial N-alkylamino acid ampholytes having application in this invention include alkyl beta-amino dipropionates, $\text{RN}(\text{C}_2\text{H}_4\text{COOM})_2$ and $\text{RNHC}_2\text{H}_4\text{COOM}$. In these R is preferably an acyclic hydrophobic group containing from about 8 to about 18 carbon atoms, and M is a cation to neutralize the charge of the anion.

Preferred amphoteric surfactants include those derived from coconut products such as coconut oil or coconut fatty acid. The more preferred of these coconut derived surfactants include as part of their structure an ethylenediamine moiety, an alkanolamide moiety, an amino acid moiety, preferably glycine, or a combination thereof; and an aliphatic substituent of from about 8 to 18 (preferably 12) carbon atoms. Such a surfactant can also be considered an alkyl amphodicarboxylic acid. These amphoteric surfactants can include chemical structures represented as: $\text{C}_{12}\text{-alkyl-C(O)-NH-CH}_2\text{-CH}_2\text{-N}^+(\text{CH}_2\text{-CH}_2\text{-CO}_2\text{Na})_2\text{-CH}_2\text{-CH}_2\text{-OH}$ or $\text{C}_{12}\text{-alkyl-C(O)-N(H)-CH}_2\text{-CH}_2\text{-N}^+(\text{CH}_2\text{-CO}_2\text{Na})_2\text{-CH}_2\text{-CH}_2\text{-OH}$. Disodium cocoampho dipropionate is one most preferred amphoteric surfactant and is commercially available under the tradename Miranol™ FBS from Rhodia Inc., Cranbury, N.J. Another most preferred coconut derived amphoteric surfactant with the chemical name disodium cocoampho diacetate is sold under the tradename Miranol™ C2M-SF Conc., also from Rhodia Inc., Cranbury, N.J.


A typical listing of amphoteric classes, and species of these surfactants, is given in U.S. Pat. No. 3,929,678 issued to Laughlin and Heuring on Dec. 30, 1975.

Further examples are given in "Surface Active Agents and Detergents" (Vol. I and II by Schwartz, Perry and Berch).

Zwitterionic Surfactants

5 Zwitterionic surfactants can be thought of as a subset of the amphoteric surfactants. Zwitterionic surfactants can be broadly described as derivatives of secondary and tertiary amines, derivatives of heterocyclic secondary and tertiary amines, or derivatives of quaternary ammonium, quaternary phosphonium or tertiary sulfonium compounds. Typically, a zwitterionic surfactant includes a positive
10 charged quaternary ammonium or, in some cases, a sulfonium or phosphonium ion; a negative charged carboxyl group; and an alkyl group. Zwitterionics generally contain cationic and anionic groups which ionize to a nearly equal degree in the isoelectric region of the molecule and which can develop strong "inner-salt" attraction between positive-negative charge centers. Examples of such zwitterionic
15 synthetic surfactants include derivatives of aliphatic quaternary ammonium, phosphonium, and sulfonium compounds, in which the aliphatic radicals can be straight chain or branched, and wherein one of the aliphatic substituents contains from 8 to 18 carbon atoms and one contains an anionic water solubilizing group, e.g., carboxy, sulfonate, sulfate, phosphate, or phosphonate. Betaine and sultaine
20 surfactants are exemplary zwitterionic surfactants for use herein.

A general formula for these compounds is:




25 wherein R¹ contains an alkyl, alkenyl, or hydroxyalkyl radical of from 8 to 18 carbon atoms having from 0 to 10 ethylene oxide moieties and from 0 to 1 glyceryl moiety; Y is selected from the group consisting of nitrogen, phosphorus, and sulfur atoms; R² is an alkyl or monohydroxy alkyl group containing 1 to 3 carbon atoms; x is 1 when Y is a sulfur atom and 2 when Y is a nitrogen or phosphorus atom, R³ is an

alkylene or hydroxy alkylene or hydroxy alkylene of from 1 to 4 carbon atoms and Z is a radical selected from the group consisting of carboxylate, sulfonate, sulfate, phosphonate, and phosphate groups.

Examples of zwitterionic surfactants having the structures listed above include: 4-[N,N-di(2-hydroxyethyl)-N-octadecylammonio]-butane-1-carboxylate; 5-[S-3-hydroxypropyl-S-hexadecylsulfonio]-3-hydroxypentane-1-sulfate; 3-[P,P-diethyl-P-3,6,9-trioxatetracosanephosphonio]-2-hydroxypropane-1-phosphate; 3-[N,N-dipropyl-N-3-dodecoxy-2-hydroxypropyl-ammonio]-propane-1-phosphonate; 3-(N,N-dimethyl-N-hexadecylammonio)-propane-1-sulfonate; 3-(N,N-dimethyl-N-hexadecylammonio)-2-hydroxy-propane-1-sulfonate; 4-[N,N-di(2(2-hydroxyethyl)-N(2-hydroxydodecyl)ammonio)-butane-1-carboxylate; 3-[S-ethyl-S-(3-dodecoxy-2-hydroxypropyl)sulfonio]-propane-1-phosphate; 3-[P,P-dimethyl-P-dodecylphosphonio]-propane-1-phosphonate; and S[N,N-di(3-hydroxypropyl)-N-hexadecylammonio]-2-hydroxy-pentane-1-sulfate. The alkyl groups contained in said detergent surfactants can be straight or branched and saturated or unsaturated.

The zwitterionic surfactant suitable for use in the present compositions includes a betaine of the general structure:



20

These surfactant betaines typically do not exhibit strong cationic or anionic characters at pH extremes nor do they show reduced water solubility in their isoelectric range. Unlike "external" quaternary ammonium salts, betaines are compatible with anionics. Examples of suitable betaines include coconut acylamidopropyldimethyl betaine; hexadecyl dimethyl betaine; C₁₂₋₁₄ acylamidopropylbetaine; C₈₋₁₄ acylamidohexyldiethyl betaine; 4-C₁₄₋₁₆ acylmethyamidodiethylammonio-1-carboxybutane; C₁₆₋₁₈

25

acylamidodimethylbetaine; C₁₂₋₁₆ acylamidopentanedithylbetaine; and C₁₂₋₁₆ acylmethylamidodimethylbetaine.

Sultaines useful in the present invention include those compounds having the formula $(R(R^1)_2 N^+ R^2 SO_3^-)$, in which R is a C₆-C₁₈ hydrocarbyl group, each R¹ is typically independently C₁-C₃ alkyl, e.g. methyl, and R² is a C₁-C₆ hydrocarbyl group, e.g. a C₁-C₃ alkylene or hydroxyalkylene group.

A typical listing of zwitterionic classes, and species of these surfactants, is given in U.S. Pat. No. 3,929,678 issued to Laughlin and Heuring on Dec. 30, 1975. Further examples are given in "Surface Active Agents and Detergents" (Vol. I and II by Schwartz, Perry and Berch).

Surfactant Compositions

The surfactants described hereinabove can be used singly or in combination in the practice and utility of the present invention. In particular, the nonionics and anionics can be used in combination. The semi-polar nonionic, cationic, amphoteric and zwitterionic surfactants can be employed in combination with nonionics or anionics. The above examples are merely specific illustrations of the numerous surfactants which can find application within the scope of this invention. The foregoing organic surfactant compounds can be formulated into any of the several commercially desirable composition forms of this invention having disclosed utility. Said compositions are washing or presoak treatments for food or other soiled surfaces in concentrated form which, when dispensed or dissolved in water, properly diluted by a proportionating device, and delivered to the target surfaces as a solution, gel or foam will provide cleaning. Said cleaning treatments consisting of one product; or, involving a two product system wherein proportions of each are utilized. Said product is typically a concentrate of liquid or emulsion.

Additional Ingredients

The present stabilized enzyme cleaning composition can include any of a variety of ingredients typically included in enzyme or other cleaning compositions.

Such ingredients include, but are not limited to, builder, divalent ion, polyol, dye, carbohydrate, and the like.

Builder

5 Detergent builders can optionally be included in the stabilized enzyme cleaning compositions of the present invention for purposes including assisting in controlling mineral hardness. Inorganic as well as organic builders can be used. The level of builder can vary widely depending upon the end use of the composition and its desired physical form. When present, the compositions will typically include
10 at least 1%, preferably about 1% to about 10%, preferably about 2% to about 6%, more preferably about 4% to about 7% by weight builder.

Inorganic or phosphate-containing detergent builders include alkali metal, ammonium and alkanolammonium salts of polyphosphates (e.g. tripolyphosphates, pyrophosphates, and glassy polymeric meta-phosphates). Non-phosphate builders
15 may also be used. These can include phytic acid, silicates, alkali metal carbonates (e.g. carbonates, bicarbonates, and sesquicarbonates), sulphates, aluminosilicates, monomeric polycarboxylates, homo or copolymeric polycarboxylic acids or their salts in which the polycarboxylic acid includes at least two carboxylic radicals separated from each other by not more than two carbon atoms, citrates, succinates,
20 and the like. Preferred builders include citrate builders, e.g., citric acid and soluble salts thereof, due to their ability to enhance detergency of a soap or detergent solution and their availability from renewable resources and their biodegradability.

Divalent Ion

25 The stabilized enzyme cleaning compositions of the invention can contain a divalent ion, selected from calcium and magnesium ions, at a level of from 0.05% to 5% by weight, preferably from 0.1% to 1% by weight, more preferably about 0.25% by weight of the composition. The divalent ion can be, for example, calcium or magnesium. Calcium ions can preferably be included in the present stabilized
30 enzyme cleaning compositions. The calcium ions can, for example, be added as a chloride, hydroxide, oxide, formate or acetate, or nitrate, preferably chloride, salt.

Polyol

The stabilized enzyme cleaning composition of the invention can also include a polyol. The polyol advantageously provides additional stability and hydrotrophic properties to the stabilized enzyme cleaning composition. Propylene glycol and sorbitol are preferred polyols.

Dye

The stabilized enzyme cleaning composition of the invention can also include a dye. The dye advantageously provides visibility of the product in a package, dispenser, and/or lines to the stabilized enzyme cleaning composition. A wide variety of dyes are suitable, including Acid Green 25 and Direct Blue 86. Preferred dyes include a dye sold under the trade name Acid Green 25.

Manual Warewashing Presoak Method

According to the manual presoaking method aspect of this invention, soiled utensils, pots, or pans are contacted with an effective amount, typically from about 0.2% to about 0.8% by weight, preferably from about 0.2% to about 0.4% by weight, of the composition of the present invention. Such an effective amount can be used to presoak, for example, about 300 utensils in about 3 to about 5 gallons of the diluted composition. The actual amount of presoak composition used will be based on the judgment of user, and will depend upon factors such as the particular product formulation of the composition, the concentration of the composition, the number of soiled articles to be presoaked and the degree of soiling of the articles. Subsequently, the items are subjected to a manual or machine washing or rinsing method, involving either further washing steps and use of detergent product, and/or to a manual or machine rinsing method.

The present invention may be better understood with reference to the following examples. These examples are intended to be representative of specific embodiments of the invention, and are not intended as limiting the scope of the invention.

EXAMPLES

Examples of stabilized enzyme cleaning compositions according to the present invention were made and the resulting enzyme stability was compared to other conventional compositions. The compositions of eight formulas that were made and compared are summarized in Table 1. The enzyme storage stability results for these compositions were determined at ambient temperature, 100 °F, and 110 °F. These results are summarized in Figures 1, 2, and 3, respectively.

10. Table 1 - - Conventional and Boric Acid Salt Enzyme Cleaning Compositions

Ingredient	#1	#2	#3	#4	#5	#6	#7	#8
Soft Water	62.98	58.98	33.30	48.73	47.73	50.23	52.73	52.73
CaCl ₂				0.25	0.25	0.25	0.25	0.25
Propylene Glycol	10.00	10.00	30.00	10.00	8.00	8.00	8.00	
Sorbitol, 70%								8.00
Miranol FBS/C2M-SF, 39 %	5.00	5.00	10.00	5.00	8.00	8.00	8.00	8.00
MEA		15.00	15.00					
KOH, 45%				20.00	20.00	17.50	15.00	15.00
Sodium Carbonate	15.00							
Boric Acid				10.00	10.00	10.00	10.00	10.00
Briquest 301-50A		9.68	9.68					
Citric Acid, Granular				4.00	4.00	4.00	4.00	4.00
Dequest 2010	5.00							
Enzyme, Purefect 4000L	2.00	2.00	2.00	2.00	2.00	2.00	2.00	2.00
Acid Green 25	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02
Total	100.00	100.68	100.00	100.00	100.00	100.00	100.00	100.00
100% pH	10.2	10.75	10.38	10	10	9.3	9.04	
.2% pH		9.82	9.47	9.34	9.27	9.13, 9.09	9.09	
Grams of Ca ²⁺ and Mg ²⁺ chelated	0.5	1.04	1.04	1	1.04	1.04, 1.00	1	1
% Water	68.03	66.97	44.29	62.73	63.53	64.66	65.78	69.13

Formula #1 provides a representative conventional composition employing ash/ATMP for maintaining an alkaline pH. As can be seen in Figures 1-3, these

formulas quickly lost their enzyme activity upon storage, even at ambient temperature.

Formulas #2 and #3 provide representative conventional compositions employing MEA/ATMP for alkalinity. Figures 1-3, illustrate that, in conventional
5 compositions, reducing water concentration to below 45% (Formula #3) increases enzyme stability compared to a composition having 67% water (Formula #2). The level of enzyme stability at 67% water is unacceptable for a commercial enzyme cleaning composition.

Formulas #4 - #8 include the boric acid salt potassium borate, which
10 maintains alkaline pH and stabilizes the enzyme. In these compositions the potassium borate was generated through the neutralization of boric acid with potassium hydroxide. Sodium borate was not sufficiently soluble to provide the concentrations achieved with potassium borate. For example, precipitate formed when sodium hydroxide was employed to neutralize boric acid at these
15 concentrations. The exact weight percent of water in Formulas #4-#8 depends on how this value is calculated. The values shown in Table 1 do not include water that might be considered to hydrate, neutralize, or conjugate to the boric acid used to make the formula. If such water is included, the values listed for weight percent of water are increased by about 2%.

20 Surprisingly, employing the boric acid salt potassium borate dramatically enhanced enzyme storage stability, even though these formulas all contain high levels of water (62.73% – 69.13%). This is illustrated in Figures 1-3. In fact, the potassium borate compositions exhibit much better enzyme stability than even Formula #3, which has much lower level of water.

25 Figures 1-3 report results obtained with a formula including a protease enzyme. As shown in Figure 1, protease in formulas of the present invention typically shows levels higher than control levels of protease. That is, the protease that has been in a liquid enzyme cleaning composition according to the invention has greater or enhanced activity compared to the same quantity of enzyme that has not
30 been in the inventive composition. The present compositions not only stabilize the enzyme, but also enhance the activity of certain enzymes, e.g. proteases.

Although not shown in the present Table or Figures, amylase enzymes were also stabilized in the liquid enzyme cleaning composition of the present invention. The amylase retained all of its initial activity upon storage at ambient temperature for at least 35 days. These results indicate that the present compositions stabilize
5 several different enzymes.

Materials

The following materials present examples of materials suitable for preparing the compositions of the present invention. Calcium Chloride: Calcium chloride
10 Pellets 90 (Dow chemical). Propylene Glycol: Propylene Glycol, Technical (Eastman Kodak, Arco Chemical, Arch Chemical, Huntsman Corporation). Sorbitol: Sorbitol solution 70% USP/FCC (Lonza, Sorini, Speciality Products Corporation, Archer Daniels Midland, Roquette Corporation). Miranol: Dicarboxylic Coconut derivative Sodium Salt, 38% (Lonza, McIntyre Group LTD,
15 Rhodia). MEA: Monoethanolamine, 99% (Dow Chemical, Huntsman Corporation, EquiStar, Union Carbide). KOH: Potassium Hydroxide, 45% (Ashta, OxyChem, Vulcan Chemical). Sodium Carbonate: Sodium Carbonate, Dense Soda Ash (North American Chemical, Vulcan, Occidental Chemical). Boric Acid: Boric Acid, Orthoboric Acid (U.S. Borax, North American Chemical). Briquest 301-50A:
20 Amino Tri (Methylene Phosphonic Acid) (ATMP), 50%, low ammonia (Albright & Wilson). Citric Acid: Citric Acid, anhydrous granular (AE Staley Mfg. Co., Huangshi Xianglung Corporation, Zhong Ya Chemical, China Huitung Corporation, Chiel Sugar). Dequest 2010: Phosphonic Acid (1-hydroxyethylidene)bis, 60% (Solutia Inc.). Purefect 4000L: Purafect 4000L, Subtilisin Protease Enzyme
25 (Genencor International). Acid Green 25: Dye, Acid Green 25 (Bayer Corporation, Crompton & Knowles).

It should be noted that, as used in this specification and the appended claims, the singular forms "a," "an," and "the" include plural referents unless the content
30 clearly dictates otherwise. Thus, for example, reference to a composition containing "a compound" includes a mixture of two or more compounds. It should also be

noted that the term “or” is generally employed in its sense including “and/or” unless the content clearly dictates otherwise.

All publications and patent applications in this specification are indicative of the level of ordinary skill in the art to which this invention pertains.

- 5 The invention has been described with reference to various specific and preferred embodiments and techniques. However, it should be understood that many variations and modifications may be made while remaining within the spirit and scope of the invention.

WE CLAIM:

1. A liquid enzyme cleaning composition comprising surfactant, deterative enzyme, boric acid salt, and at least about 60% by weight water;
5 wherein the liquid enzyme cleaning composition is formulated to provide deterative enzyme that retains about 100% of its initial activity at ambient temperature for at least about 11 months after forming the composition.
2. The composition of claim 1, wherein the boric acid salt comprises an
10 alkali metal boric acid salt, an alkanol amine boric acid salt, or a combination thereof.
3. The composition of claim 2, wherein the boric acid salt comprises monoethanolammonium borate, diethanolammonium borate, triethanolammonium
15 borate, or a combination thereof.
4. The composition of claim 1, wherein the boric acid salt comprises potassium borate.
- 20 5. The composition of claim 4, wherein the potassium borate comprises a combination of potassium hydroxide and boric acid.
6. The composition of claim 4, wherein the composition comprises about 10 to about 15 weight percent potassium borate.
25
7. The composition of claim 1, wherein the composition is a solution.
8. The composition of claim 1, wherein the composition comprises about 60% by weight to about 85% by weight water.
30
9. The composition of claim 8, wherein the composition comprises

about 60% by weight to about 70% by weight water.

10. The composition of claim 1, wherein the deterative enzyme retains at least about 80% of its initial activity at 100 °F for at least about 70 days after forming the composition.

11. The composition of claim 1, wherein the deterative enzyme retains at least about 40% of its initial activity at 120 °F for at least about 25 days after forming the composition.

10

12. The composition of claim 1, wherein the deterative enzyme comprises protease, amylase, lipase, cellulase, peroxidase, gluconase, or a combination thereof.

13. The composition of claim 12, wherein the deterative enzyme comprises alkaline protease, lipase, amylase, or a combination thereof.

14. The composition of claim 1, wherein the surfactant comprises amphoteric surfactant.

15. The composition of claim 14, wherein the amphoteric surfactant comprises a coconut derived surfactant comprising an ethylenediamine moiety, an amide moiety, an amino acid moiety, or a combination thereof; and an aliphatic moiety.

16. The composition of claim 14, wherein the amphoteric surfactant comprises an alkyl amphodicarboxylic acid.

17. The composition of claim 14, wherein the amphoteric surfactant comprises C_{12} -alkyl-C(O)-NH-CH₂-CH₂-N⁺(CH₂-CH₂-CO₂Na)₂-CH₂-CH₂-OH or C_{12} -alkyl-C(O)-N(H)-CH₂-CH₂-N⁺(CH₂-CO₂Na)₂-CH₂-CH₂-OH, or a combination thereof.

18. The composition of claim 14, wherein the amphoteric surfactant comprises disodium cocoampho dipropionate, disodium cocoampho diacetate, or a combination thereof.

5

19. The composition of claim 18, wherein the alkyl amphodicarboxylic acid surfactant comprises a surfactant sold under the trade name Miranol FBS, Miranol C2M SF conc., or a combination thereof.

10 20. The composition of claim 1, further comprising source of calcium ions, polyol, builder, dye, or a combination thereof.

21. The composition of claim 20, wherein the surfactant comprises an amphoteric surfactant, the deterative enzyme comprises protease, the boric acid salt
15 comprises potassium borate, the source of calcium ions comprises calcium chloride, the polyol comprises propylene glycol, the builder comprises citric acid salt, and the dye comprises a dye sold under the trade name Acid Green 25.

22. The composition of claim 21, comprising about 8% by weight
20 surfactant, about 2% by weight protease, about 10% to about 15% by weight potassium borate, about 0.25% by weight calcium chloride, about 8% by weight propylene glycol, about 4% to about 7% by weight citric acid salt, and about 0.02% by weight Acid Green 25.

25 23. The composition of claim 1, further comprising a pH in the range of about 9 to about 10.

24. The composition of claim 1, wherein the composition provides enhanced enzyme activity.

30

25. A liquid enzyme cleaning composition comprising surfactant,

deterative enzyme, potassium borate, and at least about 60% by weight water.

26. The composition of claim 25, wherein the potassium borate comprises a combination of potassium hydroxide and boric acid.

5

27. The composition of claim 25, wherein the composition comprises about 10 to about 15 weight percent potassium borate.

10

28. The composition of claim 25, wherein the composition is a solution.

29. The composition of claim 25, wherein the composition comprises about 60% by weight to about 85% by weight water.

15

30. The composition of claim 29, wherein the composition comprises about 60% by weight to about 70% by weight water.

20

31. The composition of claim 25, wherein the deterative enzyme retains about 100% of its initial activity at ambient temperature for at least about 11 months after forming the composition.

32. The composition of claim 25, wherein the deterative enzyme retains at least about 80% of its initial activity at 100 °F for at least about 70 days after forming the composition.

25

33. The composition of claim 25, wherein the deterative enzyme retains at least about 40% of its initial activity at 120 °F for at least about 25 days after forming the composition.

30

34. The composition of claim 25, wherein the deterative enzyme comprises protease, amylase, lipase, cellulase, peroxidase, gluconase, or a combination thereof.

35. The composition of claim 34, wherein the deterative enzyme comprises alkaline protease, lipase, amylase, or a combination thereof.

5 36. The composition of claim 25, wherein the surfactant comprises amphoteric surfactant.

37. The composition of claim 36, wherein the amphoteric surfactant comprises a coconut derived surfactant comprising an ethylenediamine moiety, an amide moiety, an amino acid moiety, or a combination thereof, and an aliphatic moiety.

38. The composition of claim 36, wherein the amphoteric surfactant comprises an alkyl amphodicarboxylic acid.

15

39. The composition of claim 36, wherein the amphoteric surfactant comprises
 C_{12} -alkyl-C(O)-NH-CH₂-CH₂-N⁺(CH₂-CH₂-CO₂Na)₂-CH₂-CH₂-OH or C_{12} -alkyl-C(O)-N(H)-CH₂-CH₂-N⁺(CH₂-CO₂Na)₂-CH₂-CH₂-OH, or a combination thereof.

20

40. The composition of claim 36, wherein the amphoteric surfactant comprises disodium cocoampho dipropionate, disodium cocoampho diacetate, or a combination thereof.

25 41. The composition of claim 40, wherein the alkyl amphodicarboxylic acid surfactant comprises a surfactant sold under the trade name Miranol FBS, Miranol C2M SF conc., or a combination thereof.

42. The composition of claim 25, further comprising source of calcium ions, polyol, builder, dye, or a combination thereof.

30

43. The composition of claim 42, wherein the surfactant comprises an amphoteric surfactant, the detergent enzyme comprises protease, the boric acid salt comprises potassium borate, the source of calcium ions comprises calcium chloride, the polyol comprises propylene glycol, the builder comprises citric acid salt, and the
5 dye comprises a dye sold under the trade name Acid Green 25.

44. The composition of claim 42, comprising about 8% by weight surfactant, about 2% by weight protease, about 10% to about 15% by weight potassium borate, about 0.25% by weight calcium chloride, about 8% by weight
10 propylene glycol, about 4% to about 7% by weight citric acid salt, and about 0.02% by weight Acid Green 25.

45. The composition of claim 25, further comprising a pH in the range of about 9 to about 10.
15

46. The composition of claim 25, wherein the composition provides enhanced enzyme activity.

47. A liquid enzyme cleaning composition comprising a surfactant, a
20 detergent enzyme, an alkanol amine boric acid salt, and at least about 60% by weight water.

48. The composition of claim 47, wherein the boric acid salt comprises monoethanolammonium borate, diethanolammonium borate, triethanolammonium
25 borate, or a combination thereof.

49. A liquid enzyme cleaning composition comprising a surfactant; a detergent enzyme; at least about 60% by weight water; and potassium borate, monoethanolammonium borate, diethanolammonium borate, triethanolammonium
30 borate, or a combination thereof.

50. A liquid enzyme cleaning composition comprising surfactant, deterative enzyme, boric acid salt, and greater than 80% by weight water.


FIG. 1

FIG. 2





FIG. 3

PROTEOLYTIC ENZYME ACTIVITY ASSAY

120°F



INTERNATIONAL SEARCH REPORT

Int. Application No.

PCT/US 01/20214

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 C11D3/386 C11D3/02

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C11D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ.

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP 0 481 542 A (UNILEVER) 22 April 1992 (1992-04-22)	1,2, 7-14,23, 24
Y	page 4, line 28 -page 7, line 28	1-4,25, 28-36, 42,46-49
A	page 8, line 7 - line 15	5,6, 15-22, 26,27, 37-45,50
	claims	
Y	WO 95 00621 A (ALLERGAN INC) 5 January 1995 (1995-01-05)	1,2,4, 25, 28-36, 42,46,49
	page 17, line 6 - line 13 page 18, line 5 - line 6	
	--- -/-- ---	



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents :

A document defining the general state of the art which is not considered to be of particular relevance

E earlier document but published on or after the international filing date

L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

O document referring to an oral disclosure, use, exhibition or other means

P document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

* & * document member of the same patent family

Date of the actual completion of the international search

17 October 2001

Date of mailing of the international search report

26/10/2001

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Serbetsoglou, A

INTERNATIONAL SEARCH REPORT

Int. Application No.

PCT/US 01/20214

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	GB 1 224 564 A (UNILEVER) 10 March 1971 (1971-03-10) page 1, line 53 - line 87 page 2, line 4 - line 18 claims 1-5 ---	1-3, 47-49
X	EP 0 451 924 A (COLGATE-PALMOLIVE CO) 16 October 1991 (1991-10-16) page 4, line 1 -page 8, line 11 page 10, line 16; claims ---	1,2, 7-14,24
A		3-6, 15-23, 25-50
X	EP 0 348 183 A (UNILEVER) 27 December 1989 (1989-12-27) page 3, line 35 -page 6, line 11 claims 1-16 ---	1,2, 7-14,20, 23,24
X	GB 2 140 818 A (COLGATE-PALMOLIVE CO) 5 December 1984 (1984-12-05) page 1, line 38 -page 4, line 36 claims 1-8 ---	1,2, 7-14,24, 50
X	GB 2 200 132 A (COLGATE-PALMOLIVE CO) 27 July 1988 (1988-07-27) page 13, line 9 - line 26 page 19, line 33 -page 20, line 10 claims 1-6 ---	1,2,4, 7-14,24
X	US 4 462 922 A (BOSKAMP JELLES V) 31 July 1984 (1984-07-31) column 1, line 50 -column 4, line 13 claims ---	1,2,4, 7-14,24
X	US 4 670 179 A (INAMORATO JACK T ET AL) 2 June 1987 (1987-06-02) column 3, line 4 -column 5, line 12 column 6, line 15 - line 17 claims 1-8 -----	1,2, 7-14,24

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 01/20214

Patent document cited in search report		Publication date	Patent family member(s)		Publication date
EP 0481542	A	22-04-1992	AU	8551991 A	09-04-1992
			BR	9104258 A	02-06-1992
			CA	2052601 A1	05-04-1992
			EP	0481542 A2	22-04-1992
			JP	6346089 A	20-12-1994
			ZA	9107956 A	05-04-1993

WO 9500621	A	05-01-1995	AT	205245 T	15-09-2001
			AU	697386 B2	01-10-1998
			AU	7208794 A	17-01-1995
			CA	2165074 A1	05-01-1995
			DE	69428202 D1	11-10-2001
			EP	0703968 A1	03-04-1996
			HU	72713 A2	28-05-1996
			IL	109705 A	15-07-1998
			JP	8511882 T	10-12-1996
			US	6165954 A	26-12-2000
			WO	9500621 A1	05-01-1995
			US	5630884 A	20-05-1997
			US	5783532 A	21-07-1998
			US	5746838 A	05-05-1998
			ZA	9404287 A	13-02-1995

GB 1224564	A	10-03-1971	AT	294284 B	15-10-1971
			BE	732312 A	29-10-1969
			CH	510733 A	31-07-1971
			DE	1921440 A1	13-11-1969
			ES	366571 A1	16-03-1971
			FR	2014122 A5	17-04-1970
			LU	58496 A1	13-11-1969
			NL	6906496 A	31-10-1969

EP 0451924	A	16-10-1991	AT	176682 T	15-02-1999
			AU	651090 B2	14-07-1994
			AU	7400791 A	17-10-1991
			BR	9101486 A	03-12-1991
			CA	2040202 A1	14-10-1991
			DE	69130880 D1	25-03-1999
			DK	451924 T3	20-09-1999
			EP	0451924 A2	16-10-1991
			ES	2130131 T3	01-07-1999
			FI	911800 A	14-10-1991
			GR	91100154 A ,B	30-07-1992
			HK	1014026 A1	19-05-2000
			IE	911245 A1	23-10-1991
			MX	173751 B	25-03-1994
			NO	911448 A	14-10-1991
			NZ	237570 A	27-09-1993
			PT	97325 A	31-01-1992
			TR	25788 A	01-09-1993
			US	5364553 A	15-11-1994
			US	5221495 A	22-06-1993
			ZA	9102598 A	30-12-1992

EP 0348183	A	27-12-1989	AU	3667189 A	04-01-1990
			BR	8903088 A	06-02-1990
			EP	0348183 A2	27-12-1989
			JP	2045599 A	15-02-1990

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 01/20214

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0348183	A	US H1776 H ZA 8904785 A	05-01-1999 27-02-1991
GB 2140818	A	05-12-1984	AT 394573 B 11-05-1992 AT 178484 A 15-10-1991 AU 564917 B2 03-09-1987 AU 2831584 A 06-12-1984 BE 899779 A1 29-11-1984 BR 8402594 A 23-04-1985 CA 1220742 A1 21-04-1987 CH 659478 A5 30-01-1987 DE 3418295 A1 06-12-1984 DK 269184 A ,B, 01-12-1984 ES 532929 D0 16-06-1987 ES 8706808 A1 16-09-1987 FR 2546901 A1 07-12-1984 GR 81585 A1 11-12-1984 HK 18392 A 13-03-1992 IT 1177763 B 26-08-1987 LU 85391 A1 21-03-1985 MX 161670 A 10-12-1990 NL 8401741 A 17-12-1984 NO 842164 A 03-12-1984 NZ 208157 A 12-11-1986 PH 20171 A 14-10-1986 PT 78654 A ,B 01-06-1984 SE 8402870 A 01-12-1984 ZA 8403666 A 24-12-1985
GB 2200132	A	27-07-1988	AU 610100 B2 16-05-1991 AU 8214787 A 16-06-1988 BE 1000341 A3 25-10-1988 CH 676006 A5 30-11-1990 DE 3741617 A1 07-07-1988 DK 658287 A 16-06-1988 ES 2007760 A6 01-07-1989 FI 875464 A ,B, 16-06-1988 FR 2608168 A1 17-06-1988 GR 871905 A1 28-04-1988 IT 1224261 B 04-10-1990 JP 63186800 A 02-08-1988 LU 87075 A1 14-07-1988 MX 169200 B 24-06-1993 NL 8703029 A 01-07-1988 NO 875195 A ,B, 16-06-1988 NZ 222833 A 26-09-1990 PT 86360 A ,B 01-01-1988 SE 8704943 A 16-06-1988 ZA 8709188 A 29-11-1989
US 4462922	A	31-07-1984	AR 228215 A1 31-01-1983 AU 555411 B2 25-09-1986 AU 9046682 A 26-05-1983 BR 8206658 A 04-10-1983 CA 1190494 A1 16-07-1985 DE 3272362 D1 04-09-1986 EP 0080223 A2 01-06-1983 JP 1350556 C 28-11-1986

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 01/20214

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 4462922	A	JP 58091800 A	31-05-1983
		JP 61016796 B	02-05-1986
		NZ 202484 A	31-05-1985
		PH 18253 A	13-05-1985
		ZA 8208331 A	27-06-1984
<hr/>			
US 4670179	A	02-06-1987	NONE
<hr/>			